

Universal Cause of High- T_c Superconductivity and Anomalous Behavior of Heavy Fermion Metals

V.R. Shaginyan ^{*,1,2,3} M.Ya. Amusia,^{3,4} A.Z. Msezane,² and K.G. Popov⁵

¹*Petersburg Nuclear Physics Institute, Russian Academy of Sciences, Gatchina, 188300, Russia*

²*CTSPS, Clark Atlanta University, Atlanta, Georgia 30314, USA*

³*Racah Institute of Physics, the Hebrew University, Jerusalem 91904, Israel*

⁴*A.F. Ioffe Physical-Technical Institute,*

Russian Academy of Sciences, St. Petersburg, 194021, Russia

⁵*Komi Science Center, Ural Division, Russian Academy of Sciences, Syktyvkar, 167982, Russia*

* E-mail: vrshag@thd.pnpi.spb.ru

Abstract

Unusual properties of strongly correlated liquid observed in the high- T_c superconductors and heavy-fermion (HF) metals are determined by quantum phase transitions taking place at their critical points. Therefore, direct experimental studies of these transitions and critical points are of crucial importance for understanding the physics of high- T_c superconductors and HF metals. In case of high- T_c superconductors such direct experimental studies are absent since at low temperatures corresponding critical points are occupied by the superconductivity. Recent experimental data on the behavior of HF metals illuminate both the nature of these critical points and the nature of the phase transitions. We show that it is of crucial importance to simultaneously carry out studies of both the high- T_c superconductivity and the anomalous behavior of HF metals. The understanding of this fact has been problematic largely because of the absence of theoretical guidance. The main features of the fermion condensation quantum phase transition (FCQPT), which are distinctive in several aspects from that of conventional quantum phase transition (CQPT), are considered. Our paper deals with these fundamental problems through studies of the behavior of quasiparticles, leading to good quantitative agreement with experimental facts. We show that in contrast to CQPT, whose physics in the critical region is dominated by thermal and quantum fluctuations and characterized by the absence of quasiparticles, the physics of a Fermi system near FCQPT or undergone FCQPT is controlled by the system of Landau-type quasiparticles. However, contrary to the conventional Landau quasiparticles, the effective mass of these strongly depends on the temperature T , magnetic fields B , the number density x , etc. Our general consideration suggests that FCQPT and the emergence of novel quasiparticles near and behind FCQPT are distinctive features of strongly correlated substances such as the high- T_c superconductors and HF metals. We show that the main properties and universal behavior of the high- T_c superconductors and HF metals can be understood within the framework of presented here theory based on FCQPT. A large number of the experimental evidences in favor of the existence of FCQPT in high- T_c superconductors and HF metals is presented. We demonstrate that the essence of strongly correlated electron liquids can be controlled by both magnetic field B and temperature T . Thus, the main properties of heavy-fermion metal such as magnetoresistance, resistivity, specific heat, magnetization, volume thermal expansion, etc, are determined by its position on the $B - T$ phase diagram. The obtained results are in good agreement with recent facts and observations.

PACS numbers: 71.27.+a, 74.20.Fg, 74.25.Jb

1. INTRODUCTION

In the last two decades a new class of materials such as heavy-fermion (HF) metals and high- T_c superconductors has been found, which display a dazzling variety of physical phenomena, see e.g. [1, 2]. The high- T_c superconductors are strongly correlated metals with normal state properties that are not at all those of a normal Landau Fermi liquid (LFL). In the case of HF metals, the electronic strong correlations result in a renormalization of the effective mass of the quasiparticles, which can exceed the bare mass by a factor up to 1000 or even diverge. These non-Fermi-liquid (NFL) systems demonstrate anomalous pure power-law temperature dependences in their low temperature properties over broad temperature ranges. It is believed that in this class the basic assumption of the LFL theory that at low energies the electrons in a metal should behave as weakly interacting quasiparticles is violated. Therefore, it is generally accepted that the fundamental physics that gives rise to the high- T_c superconductivity and NFL behavior with a recovery of the LFL behavior under the application of magnetic fields observed in HF metals and high- T_c compounds is controlled by quantum phase transitions. This has made quantum phase transitions a subject of intense current interest, see e.g. [2, 3, 4, 5].

A quantum phase transition is driven by control parameters such as the composition, number density x or magnetic fields B , and takes place at a quantum critical point (QCP) when temperature $T = 0$. QCP separates an ordered phase generated by quantum phase transition from a disordered phase. It is expected that the universal behavior is only observable if the system in question is very near QCP, for example, when the correlation length is much larger than microscopic length scales. Quantum phase transitions of this sort are quite common, and we shall call them as conventional quantum phase transitions (CQPT). In the case of CQPT, the physics is dominated by thermal and quantum fluctuations of the critical state, which is characterized by the absence of quasiparticles. It is believed that the absence of quasiparticle-like excitations is the main cause of the NFL behavior and other types of critical behavior in the quantum critical region. On the base of scaling related to the divergence of the correlation length, one can construct the critical contribution to the free energy and evaluate the corresponding properties such as critical exponents, the NFL behavior, etc. [2, 3, 4, 5]. However along this way one may expect difficulties. For example, having the only critical contribution, one has to describe different types of the behavior exhibited by different HF metals, see e.g. [6, 7, 8]. Note that HF metals are three-dimensional structures, (see e.g. [6, 9, 10]) and, thus, the type of behavior cannot be related to the dimension. The critical behavior observed in

measurements on HF metals takes place up to rather high temperatures comparable with the effective Fermi temperature T_k . For example, the thermal expansion coefficient $\alpha(T)$ measured on CeNi_2Ge_2 shows a $1/\sqrt{T}$ divergence over more than two orders of magnitude in temperature drop from 6 K down to at least 50 mK [6]. It is hardly possible to understand such a behavior on the base of the assumption of scaling when the correlation length has to be much larger than microscopic length scales. Obviously, such a situation can take place only at $T \rightarrow 0$. At some temperature $T \sim T_k$, this macroscopically large correlation length must be destroyed by thermal fluctuations.

The next problem is related to explanations of the recovery of the LFL behavior under applied magnetic fields B , observed in HF metals and the high- T_c compounds, see e.g. [1, 10, 11]. At $T \rightarrow 0$, the magnetic field dependence of the coefficient $A(B)$, causing an electrical resistivity contribution $\Delta\rho = A(B)T^2$, the Sommerfeld coefficient $\gamma(B)$ and $\chi(B)$ in specific heat, $C/T = \gamma(B)$, and magnetic susceptibility, $\chi(B)$, shows that $A(B) \sim \gamma^2(B)$ and $A(B) \sim \chi^2(B)$, so that the Kadowaki-Woods ratio, $K = A(B)/\gamma^2(B)$ [9], is B -independent and conserved [10]. Such a universal behavior is hardly possible to explain within the picture assuming the absence of quasiparticles which takes place near QCP of the corresponding CQPT. As a consequence, for example, these facts are in variance to the spin-density-wave scenario [10] and the renormalization group treatment of quantum criticality [12]. Moreover, striking recent measurements of the specific heat, charge and heat transport and the resistivity used to study the nature of magnetic field-induced QCP in heavy-fermion metal CeCoIn_5 [13, 14] certainly seem to disagree with descriptions based on CQPT.

In a system of interacting bosons at temperatures lower than the temperature of Bose-Einstein condensation [15, 16], a finite number of particles is concentrated in the lowest level. In the case of a noninteracting Bose gas at zero temperature, $T = 0$, this number is simply equal to the total number of particles in the system. In a homogeneous system of noninteracting Bosons, the lowest level is the state with zero momentum, and the ground state energy is equal to zero. For a noninteracting Fermi system such a state is impossible, and its ground state energy E_{gs} reduces to the kinetic energy and is proportional to the total number of particles. Imagine an interacting system of fermions with a pure repulsive interaction. Let us increase its interaction strength. As soon as it becomes sufficiently large and the potential energy starts to prevail over the kinetic energy, we can expect the system to undergo a phase transition at $T = 0$. This takes place when the density x tends to the critical quantum point $x \rightarrow x_{FC}$ and the kinetic energy E_k becomes frustrated, while the effective inter-electron interaction, or the Landau amplitude, being sufficiently large, starts to determine the occupation numbers of quasiparticles $n(\mathbf{p})$ which deliver the minimum value to the ground state

energy $E[n(p)]$. As a result, at $x < x_{FC}$, the function $n(\mathbf{p})$ is given by the standard equation that determines the minimum of functional $E[n(\mathbf{p})]$ [17, 18, 19, 20, 21]

$$\frac{\delta E[n(\mathbf{p})]}{\delta n(\mathbf{p})} = \mu. \quad (1.1)$$

Here we deal with three dimensional (3D) case and assume that the phase transition takes place at $x < x_{FC}$. At $T = 0$ Eq. (1.1) determines the quasiparticle distribution function $n_0(\mathbf{p})$, which delivers the minimum value to the ground state energy E . The function $n_0(\mathbf{p})$ being the signature of the new state of quantum liquids [22] does not coincide with the quasiparticle distribution function of the LFL theory in the region $(p_f - p_i)$, so that $0 < n_0(\mathbf{p}) < 1$ and $p_i < p_F < p_f$, with $p_F = (3\pi^2 x)^{1/3}$ being the Fermi momentum. Such a state was called the state with fermion condensate (FC) because quasiparticles located in the region $(p_f - p_i)$ of momentum space are pinned to the chemical potential μ [17, 19, 22]. We note that the behavior obtained for the single-particle spectrum and quasiparticle distribution functions is observed within exactly solvable models [23, 24, 25]. Lowering the potential energy, FC decreases the total energy. Unlike the Bose-Einstein condensation, which occurs even in a system of noninteracting bosons, FC can take place if the coupling constant of the interaction is large, or the corresponding Landau amplitudes are large and repulsive.

We note the remarkable peculiarity of FCQPT at $T = 0$: this transition is related to the spontaneous breaking of gauge symmetry, when the superconducting order parameter $\kappa(\mathbf{p}) = \sqrt{n_0(\mathbf{p})(1 - n_0(\mathbf{p}))}$ has a nonzero value over the region occupied by FC, with the entropy $S = 0$ [21, 26]. At small values of the pairing coupling constant λ_0 , the gap $\Delta(\mathbf{p})$ is linear in λ_0 and vanishes provided that $\lambda_0 \rightarrow 0$, while $\kappa(\mathbf{p})$ remains finite [20, 21]. As we shall see, this peculiarity allows to construct the theory of high- T_c superconductivity based on FCQPT. Thus, the state with FC cannot exist at any finite temperatures and is driven by the parameter x : at $x > x_{FC}$ the system is on the disordered side of FCQPT; at $x = x_{FC}$, Eq. (1.1) possesses the non-trivial solutions $n_0(\mathbf{p})$ with $p_i = p_F = p_f$; while at $x < x_{FC}$, the system is on the ordered side [26].

One of the most challenging problems of modern condensed matter physics is the structure and properties of Fermi systems with large coupling constants. The first solution to this problem was offered by the Landau theory of Fermi liquids, later called "normal", by introducing the notion of quasiparticles and so called amplitudes, which characterize the effective interaction among them [27]. The Landau theory can be viewed as the low energy effective theory in which high energy degrees of freedom are removed by introducing the effective amplitudes instead of strong inter-particle interaction. Usually, it is assumed that the stability of the ground state of Landau liquid is determined by

the Pomeranchuk stability conditions: the stability is violated when even one of the Landau effective interaction parameters is negative and reaches a critical value [27, 28]. Note that the new phase, at which the stability conditions are restored, can in principle be again described within the framework of the same theory. However, it has been demonstrated rather recently [17] that the Pomeranchuk stability conditions cover not all possible instabilities: one of them is missed. It corresponds to the situation when, at the temperature $T = 0$, the effective mass, the most important characteristic of Landau quasiparticles, can become infinitely large. Such a situation, leading to profound consequences, can take place when the corresponding Landau amplitude being repulsive reaches some critical value. This leads to a completely new class of strongly correlated Fermi liquids with FC [17, 19, 22], which is separated from that of a normal Fermi liquid by the fermion condensation quantum phase transition (FCQPT) [29, 30].

In the FCQPT case we are dealing with the strong coupling limit where an absolutely reliable answer cannot be given on the bases of pure theoretical first principle foundation. Therefore, the only way to verify that FC occurs is to consider both exactly solvable models and experimental facts, which can be interpreted as confirming the existence of such a state. The exactly solvable models unambiguously demonstrate that Fermi liquids with FC do exist, see e.g. [23, 24, 25]. On the other hand, these facts are seen in features of those two-dimensional (2D) systems with interacting electrons or holes, which can be represented by doped quantum wells and high- T_c superconductors. Considering the HF metals and the 2D systems of ^3He , we will show that FC exist also in these systems.

The goal of our review is to describe the behavior of Fermi systems with FC and to show that the existing data on strongly correlated liquids represented by the electronic (or hole) systems of high- T_c superconductors and HF metals can be well understood within the theory of Fermi liquids based on FCQPT. In Section 2, we review the general features of Fermi liquids with FC in their normal state. Section 3 is devoted to consideration of the superconductivity in the presence of FC. We show that the superconducting state is totally transformed by the presence of FC. For instance, the maximum value Δ_1 of the superconducting gap can be as large as $\Delta_1 \sim 0.1\varepsilon_F$, while for normal superconductors one has $\Delta_1 \sim 10^{-3}\varepsilon_F$, where ε_F is the Fermi level energy. In Section 4 we describe the quasiparticle's dispersion and its lineshape and show that they strongly deviate from the case of normal Landau liquids. In Section 5 we consider the field-induced LFL in the heavy electron liquid with FC. In Section 6 we apply our theory to explain the main properties of magnetic-field induced Landau Fermi liquid in the high- T_c metals. Section 7 is devoted to the appearance of FCQPT in

different Fermi liquids. In Section 8 we analyze the main properties of HF metals whose electronic system is placed on the disordered side of FCQPT. HF metals with the electronic system located on the ordered side of FCQPT are considered in Section 9. In Section 10 we show that FC manifests itself in the dissymmetry of tunnelling conductivity which can be observed in measurements on the high- T_c compounds and HF metals. Finally, in Section 11, we summarize our main results.

2. FERMI LIQUIDS WITH FERMION CONDENSATE

To study the universal behavior of the high- T_c superconductors and HF metals at low temperatures, we use the heavy electron liquid model in order to ignore the complications of the anisotropy of the lattice of solids and its microscopic inhomogeneity. It is possible since we consider the universal behavior demonstrated by these materials and processes related to the power-law divergences of observables such as the effective mass, specific heat, thermal expansion, etc. These divergences are determined by small momenta transferred as compared to momenta of the same order of magnitude as those of the reciprocal lattice cell, and contributions coming from them can be safely ignored. On the other side, we can simply use the common concept of the applicability of the LFL theory when describing electronic properties of metals [27]. Thus, we may usefully ignore the complications due to lattice and its anisotropy. As a result, we regard the medium as homogeneous heavy electron isotropic liquid.

2.1. Landau Theory of Fermi Liquid

Let us start by explaining the important points of the LFL theory [27]. The LFL theory rests on the notion of quasiparticles which represent elementary excitations of a Fermi liquid. Therefore these are appropriate excitations to describe the low temperature thermodynamic properties. In the case of an electron system, these are characterized by the electron's quantum numbers and effective mass M^* . The ground state energy of the system in question is a functional of the quasiparticle occupation numbers (or quasiparticle distribution function) $n(\mathbf{p}, T)$, just like the free energy $F[n(\mathbf{p}, T)]$, entropy $S[n(\mathbf{p}, T)]$, and other thermodynamic functions. From the condition that the free energy $F = E - TS$ should be minimal, we can find the distribution function

$$\frac{\delta(F - \mu N)}{\delta n(\mathbf{p}, T)} = \varepsilon(\mathbf{p}, T) - \mu(T) - T \ln \frac{1 - n(\mathbf{p}, T)}{n(\mathbf{p}, T)} = 0. \quad (2.1)$$

Here μ is the chemical potential, while

$$\varepsilon(\mathbf{p}, T) = \frac{\delta E[n(\mathbf{p}, T)]}{\delta n(\mathbf{p}, T)}, \quad (2.2)$$

is the quasiparticle energy. This energy is a functional of $n(\mathbf{p}, T)$ just like the total energy $E[n(\mathbf{p}, T)]$. The entropy $S[n(\mathbf{p}, T)]$ is given by the familiar expression [27]

$$S[n(\mathbf{p}, T)] = -2 \int [n(\mathbf{p}, T) \ln n(\mathbf{p}, T) + (1 - n(\mathbf{p}, T)) \ln(1 - n(\mathbf{p}, T))] \frac{d\mathbf{p}}{(2\pi)^3}, \quad (2.3)$$

which stems from purely combinatorial considerations. Equation (2.1) is usually presented as the Fermi-Dirac distribution

$$n(\mathbf{p}, T) = \left\{ 1 + \exp \left[\frac{(\varepsilon(\mathbf{p}, T) - \mu)}{T} \right] \right\}^{-1}. \quad (2.4)$$

At $T \rightarrow 0$, one gets from Eqs. (2.1) and (2.4) the standard solution $n(p, T \rightarrow 0) \rightarrow \theta(p_F - p)$, with $\theta(p_F - p)$ is the step function, $\varepsilon(p \simeq p_F) - \mu = p_F(p - p_F)/M_L^*$, where M_L^* is the Landau effective mass [27]

$$\frac{1}{M_L^*} = \frac{1}{p} \frac{d\varepsilon(p, T=0)}{dp} \Big|_{p=p_F}. \quad (2.5)$$

It is implied that M_L^* is positive and finite at the Fermi momentum p_F . As a result, the T -dependent corrections to M_L^* , to the quasiparticle energy $\varepsilon(\mathbf{p})$, and to other quantities, start with T^2 -terms. The effective mass is given by the well-known Landau equation

$$\frac{1}{M_L^*} = \frac{1}{M} + \sum_{\sigma_1} \int \frac{\mathbf{p}_F \mathbf{p}_1}{p_F^3} F_{\sigma, \sigma_1}(\mathbf{p}_F, \mathbf{p}_1) \frac{\partial n_{\sigma_1}(\mathbf{p}_1, T)}{\partial p_1} \frac{d\mathbf{p}_1}{(2\pi)^3}. \quad (2.6)$$

Here $F_{\sigma, \sigma_1}(\mathbf{p}_F, \mathbf{p}_1)$ is the Landau amplitude depending on the momenta $\mathbf{p}$, spins σ , and M is the bare mass of an electron. For the sake of simplicity, we omit the spin dependence of the effective mass since in the case of a homogeneous liquid and weak magnetic fields M_L^* does not noticeably depend on the spins.

Applying Eq. (2.6) at $T = 0$ and taking into account that $n(\mathbf{p}, T = 0)$ becomes the step function $\theta(p_F - p)$, we obtain the standard result

$$\frac{M_L^*}{M} = \frac{1}{1 - N_0 F^1(p_F, p_F)/3}.$$

Here N_0 is the density of states of the free Fermi gas and $F^1(p_F, p_F)$ is the p -wave component of the Landau interaction amplitude. Since in the LFL theory $x = p_F^3/3\pi^2$, the Landau amplitude can be written as $F^1(p_F, p_F) = F^1(x)$. Assume that at some critical point x_{FC} the denominator

$(1 - N_0 F^1(p_F, p_F)/3)$ tends to zero, that is $(1 - N_0 F^1(x)/3) \propto (x - x_{FC}) + a(x - x_{FC})^2 + \dots \rightarrow 0$. As a result, one obtains that $M_L^*(x)$ behaves as [31, 32]

$$\frac{M_L^*(x)}{M} \simeq A + \frac{B}{x - x_{FC}} \propto \frac{1}{r}. \quad (2.7)$$

Here A and B are constants and $r = (x - x_{FC})$ is the “distance” from the QCP of FCQPT taking place at x_{FC} . The observed behavior is in good agreement with recent experimental observations (see e.g. [33, 34]), and calculations [35, 36, 37], see Section 7 as well. In the case of electronic systems, Eq. (2.7) is valid at $x > x_{FC}$ when $r > 0$ [19, 38]. Such a behavior of the effective mass can be observed in the HF metals with a quite flat, narrow conduction band, corresponding to the large effective mass $M_L^*(x \simeq x_{FC})$, with strong electron correlations and the effective Fermi temperature $T_k \sim p_F^2/M_L^*(x)$ of the order of a few Kelvin or even lower (see e.g. [1]).

2.2. Fermion condensation quantum phase transition

As we have seen above at $T = 0$ when $r = (x - x_{FC}) \rightarrow 0$, the effective mass diverges, $M_L^*(r) \rightarrow \infty$, and eventually beyond the critical point x_{FC} the distance r becomes negative making the effective mass negative, as it follows from Eq. (2.7). To escape the possibility of being in unstable and in essence meaningless states with negative values of the effective mass, the system is to undergo a quantum phase transition at the critical point $x = x_{FC}$. Because the kinetic energy near the Fermi surface is proportional to the inverse effective mass, this phase transition is triggered by the frustrated kinetic energy and can be recognized as FCQPT [29, 39]. Therefore behind the critical point x_{FC} of this transition, the quasiparticle distribution represented by the step function does not minimize the Landau functional $E[n(\mathbf{p})]$. As a result, at $x < x_{FC}$ the quasiparticle distribution is determined by the standard equation that is used to search for the minimum of the energy functional [17]

$$\frac{\delta E[n(\mathbf{p})]}{\delta n(\mathbf{p}, T=0)} = \varepsilon(\mathbf{p}) = \mu; \quad p_i \leq p \leq p_f. \quad (2.8)$$

Equation (2.8) determines the quasiparticle distribution function $n_0(\mathbf{p})$, which minimizes the ground state energy E . Being determined by Eq. (2.8), $n_0(\mathbf{p})$ does not coincide with the step function in the region $(p_f - p_i)$, so that $0 < n_0(\mathbf{p}) < 1$, while outside the region it coincides with the step function. It follows from Eq. (2.8) that the single particle spectrum or the band is completely flat over the region. Such a state was called the state with FC because quasiparticles located in the region $(p_f - p_i)$ of momentum space are pinned to the chemical potential μ [17, 19, 22]. We note that this behavior was

obtained for the band and quasiparticle distribution functions using some exactly solvable models [24, 25].




FIG. 1: The quasiparticle distribution function $n_0(p)$ and energy $\varepsilon(p)$. Since $n_0(p)$ is the solution of Eq. (2.8) it implies $n_0(p < p_i) = 1$, $n_0(p_i < p < p_f) < 1$ and $n_0(p > p_f) = 0$, while $\varepsilon(p_i < p < p_f) = \mu$. The Fermi momentum p_F obeys the condition $p_i < p_F < p_f$.

The possible solution $n_0(\mathbf{p})$ of Eq. (2.8) and the corresponding single particle spectrum $\varepsilon(\mathbf{p})$ are shown in Fig. 1.

As we shall see in Section 3 the relevant order parameter of the FC state, $\kappa(\mathbf{p}) = \sqrt{n_0(\mathbf{p})(1 - n_0(\mathbf{p}))}$, is the order parameter of the superconducting state with the infinitely small value of the superconducting gap. Therefore the entropy of this state is zero, $S(T = 0) = 0$ [19]. Thus, this state is of pure quantum nature and cannot exist at any finite temperatures. This quantum state with FC is driven by the density x : at $x > x_{FC}$ the system is on the disordered side of FCQPT; at $x = x_{FC}$, Eq. (2.8) has the non-trivial solutions $n_0(\mathbf{p})$ with $p_i = p_F = p_f$; at $x < x_{FC}$, the system is on the ordered side. We note that the solutions $n_0(\mathbf{p})$ of Eq. (2.8) can be viewed as new to the LFL theory. Indeed, at $T = 0$, the standard solution $n(\mathbf{p}, T \rightarrow 0) \rightarrow \theta(p_F - p)$ is not the only one possible. The "anomalous" solutions of Eq. (2.1) can exist because the logarithm on the right hand side of Eq. (2.1) is finite when p belongs to the region $(p_f - p_i)$ and $0 < n_0(\mathbf{p}) < 1$. Therefore in the region $T \ln[(1 - n_0(\mathbf{p}, T))/n_0(\mathbf{p}, T)]|_{T \rightarrow 0} \rightarrow 0$, and we again arrive at Eq. (2.8).

Let us assume that with the decrease of the density (or with the growth of the interaction strength) FC has just taken place. It means that $p_i \rightarrow p_f \rightarrow p_F$, and the deviation $\delta n(\mathbf{p}) = n_0(\mathbf{p}) - \theta(p_F - p)$

is small. Expanding the functional $E[n(\mathbf{p})]$ in Taylor's series with respect to $\delta n(\mathbf{p})$ and retaining the leading terms, one obtains from Eq. (2.8) the following relation

$$\mu = \varepsilon(\mathbf{p}) = \varepsilon_0(\mathbf{p}) + \int F(\mathbf{p}, \mathbf{p}_1) \delta n(\mathbf{p}_1) \frac{d\mathbf{p}_1}{(2\pi)^2} ; \quad p_i \leq p \leq p_f , \quad (2.9)$$

where $F(\mathbf{p}, \mathbf{p}_1) = \delta^2 E / \delta n(\mathbf{p}) \delta n(\mathbf{p}_1)$ is the Landau amplitude. Both quantities, the interaction and the single-particle energy $\varepsilon_0(\mathbf{p})$ are calculated at $n(\mathbf{p}) = \theta(p_F - p)$. Equation (2.9) acquires nontrivial solutions at some density $x = x_{FC}$. Thus, FCQPT takes place if the Landau amplitudes depending on the density are positive and sufficiently large, so that the potential energy is bigger than the kinetic energy. Then the transformation of the Fermi step function $n(\mathbf{p}) = \theta(p_F - p)$ into the smooth function defined by Eq. (2.9) becomes possible [17, 19]. The system in question can be considered as a strongly correlated Fermi liquid behind the point x_{FC} . It is seen from Eq. (2.9) that the FC quasiparticles form a collective state, since their energies are defined by the macroscopical number of quasiparticles within the momentum region $(p_f - p_i)$. The shape of the excitation spectra related to FC is not affected by the Landau interaction, which, generally speaking, depends on the system's properties, including the collective states, the irregularity of the composition, impurities, etc. The only thing determined by the interaction is the width of the FC region $(p_f - p_i)$ provided the interaction is sufficiently strong to produce the FC phase transition at all. Thus, we can conclude that the spectra related to FC are of a universal form, being dependent, as we shall see in Subsections 2.3 and 3.1, on the temperature and the superconducting gap. The existence of such spectra can be viewed as the characteristic feature of the "quantum protectorate" [40, 41].

2.3. The "shadow" of the fermion condensate at finite temperatures

According to Eq. (2.1), the single-particle energy $\varepsilon(\mathbf{p}, T)$ within the interval $(p_f - p_i)$ at $T \ll T_f$ is linear in T [42]. At the Fermi level, one obtains by expanding $\ln(\dots)$ in terms of $n(\mathbf{p})$

$$\varepsilon(\mathbf{p}, T) - \mu(T) = T \ln \frac{1 - n(\mathbf{p})}{n(\mathbf{p})} \simeq T \frac{1 - 2n(\mathbf{p})}{n(\mathbf{p})} \Big|_{p \simeq p_F} . \quad (2.10)$$

Here T_f is the temperature, above which FC effects become insignificant [20]

$$\frac{T_f}{\varepsilon_F} \sim \frac{p_f^2 - p_i^2}{2M\varepsilon_F} \sim \frac{\Omega_{FC}}{\Omega_F} . \quad (2.11)$$

In this formula Ω_{FC} is the FC volume, ε_F is the Fermi energy, and Ω_F is the volume of the Fermi sphere. We note that at $T \ll T_f$ the occupation numbers $n(\mathbf{p})$ are approximately independent of T ,

being given by Eq. (2.8). At finite temperatures according to Eq. (2.10), the dispersionless plateau $\varepsilon(\mathbf{p}) = \mu$ is slightly turned counter-clockwise about μ . As a result, the plateau is just a little tilted and rounded off at the end points. According to Eqs. (2.5) and (2.10), the effective mass M_{FC}^* related to FC is given by,

$$M_{FC}^* \simeq p_F \frac{p_f - p_i}{4T} . \quad (2.12)$$

To obtain Eq. (2.12) an approximation for the derivative $dn(p)/dp \simeq -1/(p_f - p_i)$ was used. It is seen from Eq. (2.12) that at $0 < T \ll T_f$, the heavy electron liquid behaves as if it were placed at QCP, in fact it is placed at the quantum critical line $x < x_{FC}$, that is the critical behavior is observed at $T \rightarrow 0$ for all $x \leq x_{FC}$.

Having in mind that $(p_f - p_i) \ll p_F$ and using Eqs. (2.11) and (2.12), the following estimates for the effective mass M_{FC}^* are obtained:

$$\frac{M_{FC}^*}{M} \sim \frac{N(0)}{N_0(0)} \sim \frac{T_f}{T} . \quad (2.13)$$

Eqs. (2.12) and (2.13) show the temperature dependence of M_{FC}^* . In Eq. (2.13) $N_0(0)$ denotes the density of states of noninteracting electron gas, and $N(0)$ is the density of states at the Fermi level. Multiplying both sides of Eq. (2.12) by $(p_f - p_i)$, we obtain the energy scale E_0 separating the slow dispersing low energy part related to the effective mass M_{FC}^* from the faster dispersing relatively high energy part defined by the effective mass M_L^* [29, 30, 43],

$$E_0 \simeq 4T . \quad (2.14)$$

It is seen from Eq. (2.14) that the scale E_0 does not depend on the condensate volume. The single particle excitations are defined according to Eq. (2.12) by the temperature and by $(p_f - p_i)$, given by Eq. (2.8). Thus, we conclude that the one-electron spectrum is of universal form and has the features of the "quantum protectorate".

It is pertinent to note that outside the FC region the single particle spectrum is not strongly affected by the temperature, being defined by M_L^* . Thus, we come to the conclusion that a system with FC is characterized by two effective masses: M_{FC}^* which is related to the single particle spectrum at lower energy scale and M_L^* describing the spectrum at higher energy scale. The existence of two effective masses is manifested by a break (or kink) in the quasiparticle dispersion, which can be approximated by two straight lines intersecting at the energy E_0 . This break takes place at temperatures $T_c \leq T \ll T_f$, which is in accord with experimental data [44], and, as we will see, at $T \leq T_c$ which is also in accord with the experimental facts [44, 45]. Here T_c is the critical

temperature of the superconducting phase transition. The quasiparticle formalism is applicable to this problem since the width γ of single particle excitations is not large compared to their energy, being proportional to the temperature, $\gamma \sim T$ at $T > T_c$ [20]. The lineshape can be approximated by a simple Lorentzian [43]. This is consistent with experimental data obtained from scans at a constant binding energy [46] (see Sec. 4).

It is essential to have in mind, that the onset of the charge density wave instability in a many-electron system, such as an electron liquid, which takes place as soon as the effective inter-electron constant reaches its critical value $r_s = r_{cdw}$, is preceded by the unlimited growth of the effective mass, which is demonstrated Section 7. Here $r_s = r_0/a_B$ with r_0 being the average distance between electrons, while a_B is the Bohr radius. Therefore the FC occurs before the onset of the charge density wave. Hence, at $T = 0$, when r_s reaches its critical value r_{FC} corresponding to x_{FC} , $r_{FC} < r_{cdw}$, FCQPT already inevitably takes place [38]. It is pertinent to note that this growth of the effective mass with decreasing electron density was observed experimentally in a metallic 2D electron system in silicon at $r_s \simeq 7.5$ [34]. Therefore we can take this value as an estimate, $r_{FC} \sim 7.5$. On the other hand, there exist charge density waves or strong fluctuations of charge ordering in underdoped high- T_c superconductors [47, 48]. Thus, the formation of FC in high- T_c compounds can be thought more as a general property of low density electron liquid embedded in these solids rather than an unusual and anomalous solution of Eq. (2.8) [38]. Beyond the point of FCQPT, the condensate volume is proportional to $(r_s - r_{FC})$ as well as $T_f/\varepsilon_F \sim (r_s - r_{FC})/r_{FC}$ at least when $(r_s - r_{FC})/r_{FC} \ll 1$. Therefore we obtain

$$\frac{r_s - r_{FC}}{r_{FC}} \sim \frac{p_f - p_i}{p_F} \sim \frac{x_{FC} - x}{x_{FC}}. \quad (2.15)$$

FC serves as a stimulator that creates new phase transitions, which eliminates the degeneracy of the spectrum. For example FC can generate spin density waves or antiferromagnetic phase transition, thus leading to a whole variety of new properties of the system under consideration. Then, the onset of the charge density wave is preceded by FCQPT, and both of these phases can coexist at the sufficiently low density when $r_s \geq r_{cdw}$. The transition to superconductivity is strongly assisted by FC because both of the phases are characterized by the same order parameter. As a result, the superconductivity by removing the spectrum degeneracy “wins” the competition with other phase transitions up to the critical temperature T_c , see Section 3. We now turn to the consideration of the superconducting state and quasiparticle dispersions at $T \leq T_c$.

3. THE SUPERCONDUCTING STATE WITH FERMION CONDENSATE

In this Section we consider the superconducting state of 2D heavy electron liquid since the high- T_c superconductors are predominantly represented by 2D structures. On the other hand, our consideration can be easily adopted to the 3D case.

3.1. Superconducting state at $T = 0$

At $T = 0$, the ground state energy $E_{gs}[\kappa(\mathbf{p}), n(\mathbf{p})]$ of a 2D electron liquid is a functional of both the order parameter of the superconducting state $\kappa(\mathbf{p})$ and the quasiparticle occupation numbers $n(\mathbf{p})$. This energy is determined by the well-known equation of the Bardeen-Cooper-Schrieffer (BCS) weak-coupling theory of superconductivity, see e.g. [49, 50]

$$E_{gs} = E[n(\mathbf{p})] + \lambda_0 \int V(\mathbf{p}_1, \mathbf{p}_2) \kappa(\mathbf{p}_1) \kappa^*(\mathbf{p}_2) \frac{d\mathbf{p}_1 d\mathbf{p}_2}{(2\pi)^4}. \quad (3.1)$$

Here $E[n(\mathbf{p})]$ is the ground-state energy of a normal Fermi liquid, and

$$n(\mathbf{p}) = v^2(\mathbf{p}); \quad \kappa(\mathbf{p}) = v(\mathbf{p})u(\mathbf{p}) = \sqrt{n(\mathbf{p})(1 - n(\mathbf{p}))}, \quad (3.2)$$

Where $u(\mathbf{p})$ and $v(\mathbf{p})$ are the normalized coherence factors, $v^2(\mathbf{p}) + u^2(\mathbf{p}) = 1$. It is assumed that the pairing interaction $\lambda_0 V(\mathbf{p}_1, \mathbf{p}_2)$ is weak. We define the superconducting gap

$$\Delta(\mathbf{p}) = -\lambda_0 \int V(\mathbf{p}, \mathbf{p}_1) \kappa(\mathbf{p}_1) \frac{d\mathbf{p}_1}{4\pi^2}. \quad (3.3)$$

Minimizing E_{gs} with respect to $v(\mathbf{p})$ we obtain the equation connecting the single-particle energy $\varepsilon(\mathbf{p})$ to $\Delta(\mathbf{p})$,

$$\varepsilon(\mathbf{p}) - \mu = \Delta(\mathbf{p}) \frac{1 - 2v^2(\mathbf{p})}{2\kappa(\mathbf{p})}, \quad (3.4)$$

where the single-particle energy $\varepsilon(\mathbf{p})$ is determined by the Landau equation (2.2). With insertion of the value of $\kappa(\mathbf{p})$ into Eq. (3.3), we obtain the known equation for $\Delta(\mathbf{p})$

$$\Delta(\mathbf{p}) = -\frac{\lambda_0}{2} \int V(\mathbf{p}, \mathbf{p}_1) \frac{\Delta(\mathbf{p}_1)}{\sqrt{(\varepsilon(\mathbf{p}_1) - \mu)^2 + \Delta^2(\mathbf{p}_1)}} \frac{d\mathbf{p}_1}{4\pi^2}. \quad (3.5)$$

If $\lambda_0 \rightarrow 0$, then the maximum value Δ_1 of the superconducting gap $\Delta(\mathbf{p})$ tends to zero, and Eq. (3.4) reduces to Eq. (2.8)

$$\frac{\delta E[n(\mathbf{p})]}{\delta n(\mathbf{p})} = \varepsilon(\mathbf{p}) - \mu = 0, \text{ provided that } 0 < n(\mathbf{p}) < 1; \text{ or } \kappa(\mathbf{p}) \neq 0; p_i \leq p \leq p_f. \quad (3.6)$$

It is seen from Eq. (3.6) that at $x < x_{FC}$, the function $n(\mathbf{p})$ is determined by the standard equation to search the minimum of functional $E[n(\mathbf{p})]$ [17, 20]. Equation (3.6) determines the quasiparticle distribution function $n_0(\mathbf{p})$ which delivers the minimum value to the ground state energy $E_{gs}[\kappa(\mathbf{p}), n(\mathbf{p})]$ in the $\lambda_0 \rightarrow 0$ limit. Now we can study the relationships between the state defined by Eq. (3.6), or by Eq. (2.8), and the superconducting state. At $T = 0$, Eq. (3.6) defines the particular state of a Fermi-liquid with FC, for which the modulus of the order parameter $|\kappa(\mathbf{p})|$ has finite values in the range of momenta $p_i \leq p \leq p_f$, while $\Delta_1 \rightarrow 0$. Such a state can be considered as superconducting, with an infinitely small value of Δ_1 , so that the entropy of this state is equal to zero. It is obvious that this state, being driven by the quantum phase transition, disappears at $T > 0$ [29, 30]. Any quantum phase transition, which takes place at temperature $T = 0$, is determined by a control parameter other than the temperature, for instance, by pressure, magnetic field, or the number density of mobile charge carriers x . In the case of FCQPT, as we have shown in Section 2, the control parameter is the density x of the system, which determines the strength of the Landau amplitudes. FCQPT occurs at the quantum critical point $x = x_{FC}$.

As any phase transition, FCQPT is related to the order parameter, which induces a broken symmetry. As it follows from Eq. (3.2), the order parameter of the state with FC is $\kappa(\mathbf{p})$. Thus, the solutions $n_0(\mathbf{p})$ of Eq. (3.6) represent a new class of the solutions of both BCS equations and LFL equations. In contrast to the conventional BCS solutions [49], the new ones are characterized by the infinitesimal value of the superconducting gap, $\Delta_1 \rightarrow 0$, while the order parameter $\kappa(\mathbf{p})$ is finite. At the same time, in contrast to the standard solutions of the LFL theory, the new ones are characterized by the superconducting order parameter $\kappa(\mathbf{p})$, that is at $T \rightarrow 0$, the quasiparticle distribution function does not tend to the step function $\theta(p_f - p)$, being the solution of Eq. (3.6). Thus, we can conclude that the solutions of Eq. (3.6) can be viewed as common solutions of both BCS and LFL equations, while Eq. (3.6) can be derived starting from either BCS or LFL theory. As we shall see, on the basis of the peculiarities of this new class of solutions, we can define the notion of the strongly correlated Fermi liquid and explain the main properties of the high- T_c superconductivity and HF metals. It is essential that the existence of state with FC can be revealed experimentally since the order parameter $\kappa(\mathbf{p})$ is suppressed by a magnetic field B . Destroying the state with FC, the magnetic field B converts the strongly correlated Fermi liquid into the normal LFL. In that case, the magnetic field plays the role of control parameter.

When $p_i \rightarrow p_F \rightarrow p_f$, Eq. (3.6) determines the critical density x_{FC} at which FCQPT takes place. When $x < x_{FC}$, the system becomes divided into two quasiparticle subsystems: the first subsystem is

in the $(p_f - p_i)$ range and is characterized by the quasiparticles with the effective mass $M_{FC}^* \propto 1/\Delta_1$, as it follows from Eq. (3.4), while the second one is occupied by quasiparticles with finite mass M_L^* and momenta $p < p_i$. When $\lambda_0 \rightarrow 0$, the density of states near the Fermi level tends to infinity, $N(0) \sim M_{FC}^* \sim 1/\Delta_1$. The quasiparticles with M_{FC}^* occupy the same energy level and form pairs with binding energy of the order of Δ_1 and with average momentum p_0 , $p_0/p_F \sim (p_f - p_i)/p_F \ll 1$. Therefore, this state strongly resembles the Bose-Einstein condensate, in which quasiparticles occupy the same energy level. But the FC quasiparticles have to be spread over the range $(p_f - p_i)$ in momentum space due to the exclusion principle. In contrast to the Bose-Einstein condensation, the fermion condensation temperature is $T_c = 0$. And in contrast to the ordinary superconductivity, FC is formed due to the Landau repulsive interaction $F(\mathbf{p}, \mathbf{p}_1)$ rather than by the relatively weak attractive quasiparticle-quasiparticle pairing interaction $\lambda_0 V(\mathbf{p}_1, \mathbf{p}_2)$.

If $\lambda_0 \neq 0$, Δ_1 becomes finite, leading to a finite value of the effective mass M_{FC}^* , which can be obtained from Eq. (3.4) upon differentiating both parts of this equation with respect to the momentum p and using Eq. (2.5) [29, 30, 43]

$$M_{FC}^* \simeq p_F \frac{p_f - p_i}{2\Delta_1} . \quad (3.7)$$

As to the energy scale, it is determined by the parameter E_0 :

$$E_0 = \varepsilon(\mathbf{p}_f) - \varepsilon(\mathbf{p}_i) \simeq p_F \frac{(p_f - p_i)}{M_{FC}^*} \simeq 2\Delta_1 . \quad (3.8)$$

3.2. Superconducting state at finite temperatures

Let us assume that the range of FC is small, that is $(p_f - p_i)/p_F \ll 1$, and $2\Delta_1 \ll T_f$ so that the order parameter $\kappa(\mathbf{p})$ is governed mainly by FC [29, 30]. To solve Eq. (3.5) analytically, we take the BCS approximation for the interaction [49]: $\lambda_0 V(\mathbf{p}, \mathbf{p}_1) = -\lambda_0$ if $|\varepsilon(\mathbf{p}) - \mu| \leq \omega_D$, i.e. the interaction is zero outside this region, with ω_D being the characteristic energy, e.g. that of phonon. As a result, the gap becomes dependent only on the temperature, $\Delta(\mathbf{p}) = \Delta_1(T)$, being independent of the momentum, and Eq. (3.5) takes the form

$$1 = N_{FC} \lambda_0 \int_0^{E_0/2} \frac{d\xi}{\sqrt{\xi^2 + \Delta_1^2(0)}} + N_L \lambda_0 \int_{E_0/2}^{\omega_D} \frac{d\xi}{\sqrt{\xi^2 + \Delta_1^2(0)}} . \quad (3.9)$$

Here we set $\xi = \varepsilon(\mathbf{p}) - \mu$ and introduce the density of states N_{FC} in the $(p_f - p_i)$ range, or E_0 range. It follows from Eq. (3.7), that $N_{FC} = (p_f - p_i)p_F/2\pi\Delta_1(0)$. The density of states N_L in the range

$(\omega_D - E_0/2)$ has the standard form $N_L = M_L^*/2\pi$. If the energy scale $E_0 \rightarrow 0$, Eq. (3.9) reduces to the BCS equation. On the other hand, assuming that $E_0 \leq 2\omega_D$ and omitting the second integral on the right hand side of Eq. (3.9), we obtain

$$\Delta_1(0) = \frac{\lambda_0 p_F (p_f - p_F)}{2\pi} \ln(1 + \sqrt{2}) = 2\beta \varepsilon_F \frac{p_f - p_F}{p_F} \ln(1 + \sqrt{2}), \quad (3.10)$$

where the Fermi energy $\varepsilon_F = p_F^2/2M_L^*$, and the dimensionless coupling constant β is given by the relation $\beta = \lambda_0 M_L^*/2\pi$. Taking the usual values of β for conventional superconductors as $\beta \simeq 0.3$, and assuming $(p_f - p_F)/p_F \simeq 0.2$, we get from Eq. (20) a large value of $\Delta_1(0) \sim 0.1\varepsilon_F$, while for conventional superconductors one has a much smaller gap, $\Delta_1(0) \sim 10^{-3}\varepsilon_F$. Taking into account the omitted above integral, we obtain

$$\Delta_1(0) \simeq 2\beta \varepsilon_F \frac{p_f - p_F}{p_F} \ln(1 + \sqrt{2}) \left(1 + \beta \ln \frac{2\omega_D}{E_0}\right). \quad (3.11)$$

It is seen from Eq. (3.11) that the correction due to the second integral is small, provided $E_0 \simeq 2\omega_D$. Below we shall show that $2T_c \simeq \Delta_1(0)$, which leads to the conclusion that the isotope effect is absent since Δ_1 is independent of ω_D . But this effect is restored as $E_0 \rightarrow 0$.

Assuming $E_0 \sim \omega_D$ but $E_0 > \omega_D$, we see that Eq. (3.11) has no standard solutions $\Delta(p) = \Delta_1(T=0)$ because $\omega_D < \varepsilon(p \simeq p_f) - \mu$ and the interaction vanishes at these momenta. The only way to obtain solutions is to restore the condition $E_0 < \omega_D$. For instance, we can define such a momentum $p_D < p_f$ that

$$\Delta_1(0) = 2\beta \varepsilon_F \frac{p_D - p_F}{p_F} \ln(1 + \sqrt{2}) = \omega_D, \quad (3.12)$$

while the other part in the $(p_f - p_i)$ range can be occupied by a gap Δ_2 of the different sign, $\Delta_1/\Delta_2 < 0$. It follows from Eq. (3.12) that the isotope effect is preserved, while both gaps can have s -wave symmetry.

At $T \simeq T_c$, Eqs. (3.7) and (3.8) are replaced by Eqs. (2.12) and (2.14), which is valid also at $T_c \leq T \ll T_f$

$$M_{FC}^* \simeq p_F \frac{p_f - p_i}{4T_c}, \quad E_0 \simeq 4T_c; \quad \text{if } T_c \leq T \text{ then, } M_{FC}^* \simeq p_F \frac{p_f - p_i}{4T}, \quad E_0 \simeq 4T. \quad (3.13)$$

Equation (3.9) is replaced by its conventional finite temperature generalization

$$\begin{aligned} 1 = & N_{FC} \lambda_0 \int_0^{E_0/2} \frac{d\xi}{\sqrt{\xi^2 + \Delta_1^2(T)}} \tanh \frac{\sqrt{\xi^2 + \Delta_1^2(T)}}{2T} + \\ & + N_L \lambda_0 \int_{E_0/2}^{\omega_D} \frac{d\xi}{\sqrt{\xi^2 + \Delta_1^2(T)}} \tanh \frac{\sqrt{\xi^2 + \Delta_1^2(T)}}{2T}. \end{aligned} \quad (3.14)$$

Putting $\Delta_1(T \rightarrow T_c) \rightarrow 0$, we obtain from Eq. (3.14)

$$2T_c \simeq \Delta_1(0) , \quad (3.15)$$

with $\Delta_1(T = 0)$ being given by Eq. (3.11). Upon comparing Eqs. (3.7), (3.8) and (3.13), (3.15), we see that M_{FC}^* and E_0 are almost temperature independent at $T \leq T_c$.

3.3. Bogoliubov quasiparticles

It is seen from Eq. (3.5) that the superconducting gap depends on the single-particle spectrum $\varepsilon(\mathbf{p})$. On the other hand, it follows from Eq. (3.4) that $\varepsilon(\mathbf{p})$ depends on $\Delta(\mathbf{p})$, provided that at $\lambda_0 \rightarrow 0$ Eq. (3.6) has the solution determining the existence of FC. Let us assume that λ_0 is small enough, so that the particle-particle interaction $\lambda_0 V(\mathbf{p}, \mathbf{p}_1)$ can only lead to a small perturbation of the order parameter $\kappa(\mathbf{p})$ determined by Eq. (3.6). It follows from Eq. (3.7) that the effective mass and the density of states $N(0) \propto M_{FC}^* \propto 1/\Delta_1$ are finite. As a result, we conclude that in contrast to the conventional theory of superconductivity the single-particle spectrum $\varepsilon(\mathbf{p})$ strongly depends on the superconducting gap and we have to solve Eqs. (2.2) and (3.5) in a self-consistent way. On the other hand, let us assume that Eqs. (2.2) and (3.5) are solved, and the effective mass M_{FC}^* is determined. Now one can fix the quasiparticle dispersion $\varepsilon(\mathbf{p})$ by choosing the effective mass M^* of the system in question equal to the obtained M_{FC}^* and then solve Eq. (3.5) as it is done in the case of the conventional BCS theory of superconductivity [49]. As a result, one observes that the superconducting state is characterized by the Bogoliubov quasiparticles (BQ) [51] with the dispersion

$$E(\mathbf{p}) = \sqrt{(\varepsilon(\mathbf{p}) - \mu)^2 + \Delta^2(\mathbf{p})},$$

and the normalization condition for the coherence factors $v(\mathbf{p})$ and $u(\mathbf{p})$ is held. We arrive to the conclusion that the observed features agree with the behavior of BQ predicted from the BCS theory. This observation suggests that the superconducting state with FC is BCS-like and implies the basic validity of the BCS formalism in describing the superconducting state [52]. Although the maximum value of the superconducting gap given by Eq. (3.11) and other exotic properties are determined by the presence of the fermion condensate. It is exactly the case that was observed experimentally in the high- T_c cuprate $\text{Bi}_2\text{Sr}_2\text{Ca}_2\text{Cu}_3\text{O}_{10+\delta}$ [53].

We have returned back to the LFL theory since the high energy degrees of freedom are eliminated and the quasiparticles are introduced. The only difference between the Landau Fermi-liquid, which

serves as a basis when constructing the superconducting state, and Fermi liquid after FCQPT is that we have to expand the number of relevant low energy degrees of freedom by introducing a new type of quasiparticles with the effective mass M_{FC}^* given by Eq. (3.7) and the energy scale E_0 given by Eq. (3.8). Therefore, the dispersion $\varepsilon(\mathbf{p})$ is characterized by two effective masses M_L^* and M_{FC}^* and by the scale E_0 , which define the low temperature properties including the line shape of quasiparticle excitations [29, 30, 54], while the dispersion of BQ has the standard form. We note that both the effective mass M_{FC}^* and the scale E_0 are temperature independent at $T < T_c$, where T_c is the critical temperature of the superconducting phase transition [54]. At $T > T_c$, the effective mass M_{FC}^* and the scale E_0 are given by Eqs. (2.12) and (2.14) respectively. Obviously, we cannot directly relate these new Landau Fermi-liquid quasiparticle excitations with the quasiparticle excitations of an ideal Fermi gas because the system in question has undergone FCQPT. Nonetheless, the main basis of the Landau Fermi liquid theory survives FCQPT: the notion of order parameter is preserved, and the low energy excitations of the strongly correlated liquid with FC are quasiparticles.

As it was shown above, properties of these new quasiparticles are closely related to the properties of the superconducting state. We may say that the quasiparticle system in the range $(p_f - p_i)$ becomes very “soft” and is to be considered as a strongly correlated liquid. On the other hand, the system’s properties and dynamics are dominated by a strong collective effect having its origin in its proximity to FCQPT and determined by the macroscopic number of quasiparticles forming FC in the range $(p_f - p_i)$. Such a system cannot be perturbed by the scattering of individual quasiparticles and has features of a “quantum protectorate” and demonstrates the universal behavior [29, 30, 40, 41]. A few remarks related to the quantum protectorate and the universal behavior [40] are in order here. As the Landau theory of Fermi liquid, the theory of the high-temperature superconductivity based on FCQPT deals with the quasiparticles which are elementary excitations of low energy. As a result, this theory gives general qualitative description of both the superconducting and normal states. Of course, one can choose the phenomenological parameters and obtain the quantitative consideration of the superconductivity as it can be done in the framework of the Landau theory when describing a normal Fermi-liquid, say ^3He . Therefore, we conclude that any theory, which is capable to describe FC and compatible with the BCS theory, will give the qualitative picture of the superconducting state and the normal state that coincides with the picture based on FCQPT. Both of the approaches can be agreed at a numerical level provided the corresponding parameters are adjusted. For example, since the formation of the flat band is possible in the Hubbard model [25], generally speaking one can repeat the results of the theory based on FCQPT in the Hubbard model. It is appropriate to

mention here that the corresponding numerical description limited to the case of $T = 0$ has been obtained within the Hubbard model [55, 56].

3.4. Pseudogap

Now let us discuss some special features of the superconducting state with FC [57, 58]. Consider two possible types of the superconducting gap $\Delta(\mathbf{p})$, namely that given by Eq. (3.5) and defined by the interaction $\lambda_0 V(\mathbf{p}, \mathbf{p}_1)$. If this interaction is dominated by an attractive phonon-mediated attraction, the solution of Eq. (3.5) with the s -wave, or the $s + d$ mixed waves will have the lowest energy. Provided the pairing interaction $\lambda_0 V(\mathbf{p}_1, \mathbf{p}_2)$ is the combination of both the attractive interaction and sufficiently strong repulsive interaction, the d -wave superconductivity can take place (see e.g. [59, 60]). But both the s -wave symmetry and the d -wave one lead to approximately the same value of the gap Δ_1 in Eq. (3.11) [54]. Therefore the non-universal pairing symmetries in high- T_c superconductivity are likely the result of the pairing interaction, while the d -wave pairing symmetry is not essential. This point of view is supported by the data [61, 62, 63, 64, 65].

We can define the critical temperature T^* , at which the superconductivity vanishes, as the temperature when $\Delta_1(T^*) \equiv 0$. At $T \geq T^*$, Eq. (3.14) has only the trivial solution $\Delta_1 \equiv 0$. On the other hand, the critical temperature T_c can be defined as a temperature, at which the superconductivity disappears while the gap occupies only part of the Fermi surface. Thus, as we shall see there are two different temperatures T_c and T^* in the case of the d -wave symmetry of the gap. It was shown [43, 57] that in the presence of FC there exist nontrivial solutions of Eq. (3.14) at $T_c \leq T \leq T^*$ when the BCS like interaction is replaced by the pairing interaction $\lambda_0 V(\mathbf{p}_1, \mathbf{p}_2)$, which includes the strong repulsive interaction leading to d -wave superconductivity. In that case, the gap $\Delta(\mathbf{p})$ as a function of angle ϕ , $\Delta(\mathbf{p}) = \Delta(p_F, \phi)$ possesses new nodes at $T > T_{node}$, as it is illustrated by Fig. 2 [43].

We show in Fig. 2 the ratio $\Delta(p_F, \phi)/T^*$ calculated at three different temperatures $0.9 T_{node}$, T_{node} , and $1.2 T_{node}$. An important difference between curve (a) and (b) and (c) is additional variations of the curves (b) and (c) marked by the two arrows. As seen from Fig. 2, the flattening occurs due to the appearance of the two new nodes. They appear at $T = T_{node}$ and move apart with increase of the temperature confining the area θ_c shown by the two arrows in Fig. 2. It is also seen from Fig. 2 that the gap Δ is extremely small over the range θ_c . It was recently shown in a number of papers (see, e.g., [66, 67]) that there exists an interplay between the magnetism and the superconductivity order parameters, leading to the damping of the magnetism order parameter below T_c . And vice versa,




FIG. 2: The gap $\Delta(p_F, \phi)$ as a function of ϕ calculated at three different temperatures expressed in terms of $T_{node} \simeq T_c$, while Δ is presented in terms of T^* . Curve (a), solid line, shows the gap calculated at temperature $0.9T_{node}$. Curve (b), dashed line, presents the same at $T = T_{node}$, and curve (c), dotted line, shows the gap calculated at temperature $1.2T_{node}$. Note the important difference in the curves (b) and (c) compared with curve (a) due to the flattening of the curves (b) and (c) about the nodes. The two arrows indicate the area θ_c emerging at T_{node} .

one can anticipate the damping of the superconductivity order parameter by the magnetism. Thus, we conclude that the gap in the range θ_c can be destroyed by strong antiferromagnetic correlations (or by spin density waves) existing in optimally doped and underdoped superconductors. It is believed that impurities can easily destroy the gap $\Delta(\mathbf{p})$ in the considered area. As a result, the superconducting gap vanishes in the macroscopic area θ_c and causes the superconductivity to die out. We have to conclude that $T_c \simeq T_{node}$, with the exact value of T_c defined by the competition between the antiferromagnetic correlations (or spin density waves) and the superconducting correlations over the range θ_c . The behavior and the shape of the pseudogap is very similar to the ones of the superconducting gap as it is seen from Fig. 2. The main difference seen from Fig. 2 is that the pseudogap vanishes along segments θ_c of the Fermi surface, while the gap vanishes at the isolated

d -wave nodes. Our calculations show that the function $\theta_c(x)$ increases very fast at small values of x , $\theta_c(x) \simeq \sqrt{x}$. Therefore T_c has to be close to T_{node} . This result is in accord with numerical calculations of the function $\theta_c([T - T_c]/T_c)$ plotted in Fig. 3.




FIG. 3: Calculated angle θ_c , pulling apart the two nodes, as a function of $(T - T_c)/T_c$.

Thus, we observe that the pseudogap state appears at temperatures $T \geq T_c \simeq T_{node}$ and vanishes at $T \geq T^*$ when Eq. (3.14) has only the trivial solution $\Delta_1 \equiv 0$. Quite naturally, one has to recognize that Δ_1 scales with T^* , while Eq. (3.15) takes the form

$$2T^* \simeq \Delta_1(0) . \quad (3.16)$$

It can then be concluded that the temperature T^* has the physical meaning of the BCS transition temperature between the state with the order parameter $\kappa \neq 0$ and the normal state.

At $T < T_c$ quasiparticle excitations are characterized by sharp peaks. When temperature becomes $T > T_c$ and $\Delta(\theta) \equiv 0$ in the range θ_c , there appear normal quasiparticle excitations with the width γ along the segments θ_c of the Fermi surface. There exists the pseudogap outside the segments θ_c , and the Fermi level is occupied by the BCS-type excitations. Both types of excitations have the width of the same order of magnitude transferring their energy and momentum to the normal quasiparticle excitations. We now estimate γ . For the entire Fermi level occupied by the normal state, the width is equal to $\gamma \approx N(0)^3 T^2 / \varepsilon(T)^2$, with the density of states $N(0) \sim M^*(T) \sim 1/T$ (see Eq. (2.12)). The dielectric constant $\varepsilon(T) \sim N(0)$, and the width γ becomes $\gamma \sim T$ [20]. In our case, however, only a part of the Fermi surface within θ_c is occupied by the normal excitations. Therefore, the

number of states allowed for quasiparticles and for quasiholes is proportional to θ_c , and the factor T^2 is replaced by $T^2\theta_c^2$. Taking this into account, we obtain $\gamma \sim \theta_c^2 T \sim T(T - T_c)/T_c \sim (T - T_c)$. Here we have omitted the small contribution coming from the BCS-type excitations. That is why the width γ vanishes at $T = T_c$. Thus, the presented above analysis shows that in the pseudogap state at $T > T_c$, the superconducting gap smoothly transforms into the pseudogap. The excitations of that area of the Fermi surface that has the gap are of the same width $\gamma \sim (T - T_c)$. The region occupied by the pseudogap is shrinking with increasing temperature. It is worth noting that the normal state resistivity $\rho(T)$ behaves as $\rho(T) \propto T$ due to $\gamma \sim (T - T_c)$. Obviously at $T > T^*$ the behavior $\rho(T) \propto T$ remains valid up to temperatures $T \sim T_f$, and T_f can be as high as the Fermi energy, provided that FC occupies noticeable part of the Fermi volume, see Eq. (2.11).

The temperature T_{node} is determined mainly by the repulsive interaction being part of the pairing interaction $\lambda_0 V(\mathbf{p}_1, \mathbf{p}_2)$. In its turn, the repulsive interaction can depend on the properties of materials such as the composition, doping, etc. Since the superconductivity is destroyed at T_c , the ratio $2\Delta_1/T_c$ can vary in a wide range and strongly depends upon the material's properties, as it follows from considerations given above [43, 57, 58]. The ratio $2\Delta_1/T_c$ can reach very high values. For instance, in the case of $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_{6+\delta}$ where the superconductivity and the pseudogap are considered to be of the common origin, $2\Delta_1/T_c$ is about 28, while the ratio $2\Delta_1/T^* \simeq 4$, which is in agreement with the experimental data for various cuprates [59]. Note that Eq. (3.16) gives also good description of the maximum gap Δ_1 in the case of the d-wave superconductivity, because the different regions with the maximum absolute value of Δ_1 and the maximal density of states can be considered as disconnected [60]. Therefore the gap in this region is formed by the attractive phonon interaction, which is approximately independent of the momenta. We can also conclude, that in the case of the s-wave pairing the pseudogap phenomenon is absent because there is no the sufficiently strong repulsive interaction. Thus, the transition from superconducting gap to pseudogap can take place only in the case of the d-wave pairing, so that the superconductivity is destroyed at $T_c \simeq T_{node}$, with the superconducting gap being smoothly transformed into the pseudogap, which closes at some temperature $T^* > T_c$ [43, 57, 58]. In the case of the s-wave pairing, we expect the absence of the pseudogap phenomenon in accordance with the observations (see e.g. [65] and references therein).

3.5. Dependence of T_c on the doping

We now turn to consideration of the maximum value of the superconducting gap Δ_1 as a function of the number density x of the mobile charge carriers which is proportional to the doping. Using Eq. (2.15), we can rewrite Eq. (3.10) as follow

$$\frac{\Delta_1}{\varepsilon_F} \sim \beta \frac{(x_{FC} - x)x}{x_{FC}}. \quad (3.17)$$

Here we take into account that the Fermi level $\varepsilon_F \propto p_F^2$, the density $x \sim p_F^2/(2M_L^*)$, and thus, $\varepsilon_F \propto x$. We can reliably assume that $T_c \propto \Delta_1$ because the empirically obtained simple bell-shaped curve of $T_c(x)$ in the high temperature superconductors [2] should have only a smooth dependence upon x . Then, $T_c(x)$ in accordance with the data has the same bell-shaped form [68]

$$T_c(x) \propto \beta(x_{FC} - x)x. \quad (3.18)$$

As an example of the implementation of the previous analysis, let us consider the main features of a room-temperature superconductor. The superconductor has to be a quasi two-dimensional structure like cuprates. From Eq. (3.10) it follows, that $\Delta_1 \sim \beta \varepsilon_F \propto \beta/r_s^2$. Noting that FCQPT in 3D systems takes place at $r_s \sim 20$ and in 2D systems at $r_s \sim 8$ [38], we can expect that Δ_1 of 3D systems comprises 10% of the corresponding maximum value of 2D superconducting gap, reaching a value as high as 60 meV for underdoped crystals with $T_c = 70$ [69]. On the other hand, it is seen from Eq. (3.10), that Δ_1 can be even large, $\Delta_1 \sim 75$ meV, and one can expect $T_c \sim 300$ K in the case of the s -wave pairing as it follows from the simple relation $2T_c \simeq \Delta_1$. Indeed, we can safely take $\varepsilon_F \sim 300$ meV, $\beta \sim 0.5$ and $(p_f - p_i)/p_F \sim 0.5$. Thus, a possible room-temperature superconductor has to be the s -wave superconductor in order to get rid of the pseudogap phenomena, which can tremendously reduce the transition temperature T_c . The density x of the mobile charge carriers must satisfy the condition $x \leq x_{FC}$ and be adjustable to reach the optimal doping level $x_{opt} \simeq x_{FC}/2$.

3.6. The gap and specific heat near T_c

Now we turn to the calculations of the gap and the specific heat at the temperatures $T \rightarrow T_c$. It is worth noting that this consideration is valid provided $T^* = T_c$. Otherwise the considered below discontinuity in the specific heat is smoothed out over the temperature range $T^* \div T_c$. For the sake of simplicity, we calculate the main contribution to the gap and the specific heat coming from FC. The function $\Delta_1(T \rightarrow T_c)$ is found from Eq. (3.14) by expanding the right hand side of the first

integral in powers of Δ_1 and omitting the contribution from the second integral on the right hand side of Eq. (3.14). This procedure leads to the following equation [54]

$$\Delta_1(T) \simeq 3.4T_c \sqrt{1 - \frac{T}{T_c}}. \quad (3.19)$$

Thus, the gap in the spectrum of the single-particle excitations has the usual behavior. To calculate the specific heat, the conventional expression for the entropy S [49] can be used

$$S = -2 \int [f(\mathbf{p}) \ln f(\mathbf{p}) + (1 - f(\mathbf{p})) \ln(1 - f(\mathbf{p}))] \frac{d\mathbf{p}}{(2\pi)^2}, \quad (3.20)$$

where

$$f(\mathbf{p}) = \frac{1}{1 + \exp[E(\mathbf{p})/T]}; \quad E(\mathbf{p}) = \sqrt{(\varepsilon(\mathbf{p}) - \mu)^2 + \Delta_1^2(T)}. \quad (3.21)$$

The specific heat C is determined by the equation

$$\begin{aligned} C = T \frac{dS}{dT} \simeq & 4 \frac{N_{FC}}{T^2} \int_0^{E_0} f(E)(1 - f(E)) \left[E^2 + T \Delta_1(T) \frac{d\Delta_1(T)}{dT} \right] d\xi + \\ & + 4 \frac{N_L}{T^2} \int_{E_0}^{\omega_D} f(E)(1 - f(E)) \left[E^2 + T \Delta_1(T) \frac{d\Delta_1(T)}{dT} \right] d\xi. \end{aligned} \quad (3.22)$$

In deriving Eq. (3.22) we again used the variable ξ and the densities of states N_{FC} and N_L , just as before in connection with Eq. (3.9), and employed the notation $E = \sqrt{\xi^2 + \Delta_1^2(T)}$. Eq. (3.22) predicts the discontinuity $\delta C = C_s - C_n$ in the specific heat C at T_c because of the two last term in the square brackets on right hand side of Eq. (3.22). Here, C_s and C_n are the specific heat of the superconducting state and the normal one respectively. Using Eq. (3.19) to calculate the first term on the right hand side of Eq. (3.22), we obtain [54]

$$\delta C(T_c) \simeq \frac{3}{2\pi^2} (p_f - p_i) p_F^2. \quad (3.23)$$

This is in contrast to the conventional result where the discontinuity is a linear function of T_c . $\delta C(T_c)$ is independent of the critical temperature T_c because as seen from Eq. (3.13) the density of states varies inversely with T_c . Note that in deriving Eq. (3.23) we took into account the main contribution coming from FC. This term vanishes as soon as $E_0 \rightarrow 0$ and the second integral on the right hand side of Eq. (3.22) gives the conventional result.

A few remarks are in order here. As we shall demonstrate in Section 9, Eq. (9.4), the specific heat of systems with FC behaves as $C(T) \propto \sqrt{T/T_f}$. The specific heat discontinuity given by Eq. (3.23) is temperature independent. As a result, we obtain that

$$\frac{\delta C(T_c)}{C_n(T_c)} \sim \sqrt{\frac{T_f}{T_c}} \frac{(p_f - p_i)}{p_F}. \quad (3.24)$$

In contrast to the conventional case of normal superconductors, when $\delta C(T_c)/C_n(T_c) = 1.43$ [27], it is seen from Eq. (3.24) that the ratio $\delta C(T_c)/C_n(T_c)$ is not the constant and can be very large provided that $T_c/T_f \ll 1$.

4. THE DISPERSION AND LINESHAPE OF THE SINGLE-PARTICLE SPECTRA

The newly discovered additional energy scale manifests itself as a break in the quasiparticle dispersion near $(50 - 70)$ meV, which results in a drastic change of the quasiparticle velocity [44, 45, 46]. Such a behavior is qualitatively different from what one could expect in a normal Fermi liquid. Moreover, this behavior can hardly be understood in the frames of either the Marginal Fermi Liquid theory or the quantum protectorate since there are no additional energy scales, or parameters, in these theories [41, 70, 71]. One could suggest that this observed strong self-energy effect, leading to the new energy scale, is due to the electron coupling with collective excitations. But in that case one has to give up the quantum protectorate idea, which would contradict observations [40, 41].

As we have seen in Sections 2 and 3, Eqs. (2.12) and (3.7), the system with FC is characterized by two effective masses: M_{FC}^* that is related to the single particle spectrum at lower energy scale, and M_L^* describing the spectrum at higher energy scale. These two effective masses manifest itself as a break in the quasiparticle dispersion, which can be approximated by two straight lines intersecting at the energy E_0 , Eqs. (2.14) and (3.8). This break takes place at temperatures $T \ll T_f$ when the system both in its superconducting state and normal one. This behavior is in good agreement with the experimental findings [44]. It is pertinent to note that at $T < T_c$, the effective mass M_{FC}^* does not depend on the momenta p_F , p_f and p_i as it follows from Eqs. (3.7) and (3.10),

$$M_{FC}^* \sim \frac{2\pi}{\lambda_0}. \quad (4.1)$$

Thus, it is seen from Eq. (4.1) that M_{FC}^* does not depend on x . This result is in good agreement with experimental facts [72, 73, 74]. The same is true for the dependence of the Fermi velocity $v_F = p_F/M_{FC}^*$ on x , because the Fermi momentum $p_F \sim \sqrt{n}$ only slightly depends on the electron density $n = n_0(1 - x)$ [72, 73]. Here n_0 is the single-particle electron density at the half-filling.

Since λ_0 is the coupling constant defining the pairing interaction, for example phonon-electron interaction, it could be reasonable to expect that the break in the quasiparticle dispersion comes from the phonon-electron interaction. The phonon scenario could explain the persistence of the break at $T > T_c$, since phonons are T -independent. On the other hand, it was shown that the quasiparticle

dispersion tends to recover to the conventional one-electron dispersion when the energy is well above the typical phonon energies [75]. The experimental observations do not show that the recovery to the one-electron dispersion takes place [44].

The lineshape function $L(q, \omega)$ of the single-particle spectrum is a function of two variables. Measurements carried out at a fixed binding energy $\omega = \omega_0$, with ω_0 being the energy of a single-particle excitation, determine the lineshape $L(q, \omega = \omega_0)$ as a function of the momentum q . We have shown above that M_{FC}^* is finite and constant at $T \leq T_c$. Therefore, at excitation energies $\omega \leq E_0$, the system behaves like an ordinary superconducting Fermi liquid with the effective mass given by Eq. (3.7) [29, 30, 43]. At $T_c \leq T$ the low energy effective mass M_{FC}^* is finite and is given by Eq. (2.12). Once again, at the energies $\omega < E_0$, the system behaves as a Fermi liquid, the single-particle spectrum is well defined while the width of single-particle excitations is of the order of T [20, 29, 30]. This behavior was observed in experiments measuring the lineshape at a fixed energy [46, 76].

The lineshape can also be determined as a function of ω , $L(q = q_0, \omega)$, at a fixed $q = q_0$. At small ω , the lineshape resembles the considered above, and $L(q = q_0, \omega)$ has the characteristic maximum and width. At energies $\omega \geq E_0$, the contribution coming from quasiparticles with the mass M_L^* become important, leading to the increase of $L(q = q_0, \omega)$. As a result, the function $L(q = q_0, \omega)$ possesses the known peak-dip-hump structure [77] directly defined by the existence of the two effective masses M_{FC}^* and M_L^* [29, 30, 43]. We can conclude that in contrast to the Landau quasiparticles, these quasiparticles have a more complicated lineshape.

To develop deeper quantitative and analytical insight into the problem, we use the Kramers-Krönig transformation to construct the imaginary part $\text{Im}\Sigma(\mathbf{p}, \varepsilon)$ of the single-particle self-energy $\Sigma(\mathbf{p}, \varepsilon)$ starting with the real one $\text{Re}\Sigma(\mathbf{p}, \varepsilon)$, which defines the effective mass [78]

$$\frac{1}{M^*} = \left(\frac{1}{M} + \frac{1}{p_F} \frac{\partial \text{Re}\Sigma}{\partial p} \right) \bigg/ \left(1 - \frac{\partial \text{Re}\Sigma}{\partial \varepsilon} \right). \quad (4.2)$$

Here M is the bare mass, while the relevant momenta p and energies ε obey the following strong inequalities: $|p - p_F|/p_F \ll 1$, and $\varepsilon/\varepsilon_F \ll 1$. We take $\text{Re}\Sigma(\mathbf{p}, \varepsilon)$ in the simplest form which accounts for the change of the effective mass at the energy scale E_0 :

$$\text{Re} \Sigma(\mathbf{p}, \varepsilon) = -\varepsilon \frac{M_{FC}^*}{M} + \left(\varepsilon - \frac{E_0}{2} \right) \frac{M_{FC}^* - M_L^*}{M} \left[\theta \left(\varepsilon - \frac{E_0}{2} \right) + \theta \left(-\varepsilon - \frac{E_0}{2} \right) \right]. \quad (4.3)$$

Here $\theta(\varepsilon)$ is the step function. Note that in order to ensure a smooth transition from the single-particle spectrum characterized by M_{FC}^* to the spectrum defined by M_L^* the step function is to be substituted by some smooth function. Upon inserting Eq. (4.3) into Eq. (4.2) we can check that

inside the interval $(-E_0/2, E_0/2)$ the effective mass can be estimated as $M^* \simeq M_{FC}^*$, and outside the interval it is $M^* \simeq M_L^*$. By applying the Kramers-Krönig transformation to $\text{Re}\Sigma(\mathbf{p}, \varepsilon)$, we obtain the imaginary part of the self-energy [54]

$$\text{Im } \Sigma(\mathbf{p}, \varepsilon) \sim \varepsilon^2 \frac{M_{FC}^*}{\varepsilon_F M} + \frac{M_{FC}^* - M_L^*}{M} \left(\varepsilon \ln \left| \frac{\varepsilon + E_0/2}{\varepsilon - E_0/2} \right| + \frac{E_0}{2} \ln \left| \frac{\varepsilon^2 - E_0^2/4}{E_0^2/4} \right| \right). \quad (4.4)$$

We see from Eq. (4.4) that at $\varepsilon/E_0 \ll 1$ the imaginary part is proportional to ε^2 , at $2\varepsilon/E_0 \simeq 1$ $\text{Im}\Sigma \sim \varepsilon$, and at $E_0/\varepsilon \ll 1$ the main contribution to the imaginary part is approximately constant. This is the behavior that gives rise to the known peak-dip-hump structure. It is seen from Eq. (4.4) that when $E_0 \rightarrow 0$ the second term on the right hand side tends to zero and the single-particle excitations become better defined, resembling the situation in a normal Fermi-liquid, and the peak-dip-hump structure eventually vanishes. On the other hand, the so called renormalization constant, or the quasiparticle renormalization factor, $a(\mathbf{p})$ is given by [78]

$$\frac{1}{a(\mathbf{p})} = 1 - \frac{\partial \text{Re } \Sigma(\mathbf{p}, \varepsilon)}{\partial \varepsilon}. \quad (4.5)$$

At $T \leq T_c$, as seen from Eqs. (4.3) and (4.5), the quasiparticle amplitude at the Fermi surface rises as the energy scale E_0 decreases. It follows from Eqs. (3.8) and (3.18) that $E_0 \sim (x_{FC} - x)/x_{FC}$. At $T > T_c$, it is seen from Eqs. (4.3) and (4.5) that the quasiparticle amplitude rises as the effective mass M_{FC}^* decreases. As seen from Eqs. (2.12) and (2.15), $M_{FC}^* \sim (p_f - p_i)/p_F \sim (x_{FC} - x)/x_{FC}$. As a result, we can conclude that the amplitude rises as the level of doping increases, while the peak-dip-hump structure vanishes and the single-particle excitations become better defined in highly overdoped samples. At $x > x_{FC}$, the energy scale $E_0 = 0$ and the quasiparticles are normal excitations of Landau Fermi liquid. It is worth noting that such a behavior was observed experimentally in highly overdoped Bi2212 where the gap size is about 10 meV [79]. Such a small size of the gap verifies that the region occupied by the FC is small since $E_0/2 \simeq \Delta_1$. Then, recent experimental data have shown that the Landau Fermi liquid does exist in heavily overdoped non-superconducting $\text{La}_{1.7}\text{Sr}_{0.3}\text{CuO}_4$ [80, 81].

5. FIELD-INDUCED LFL IN HEAVY ELECTRON LIQUID WITH FC

In this Section we consider the behavior of the heavy-electron liquid with FC in magnetic fields, assuming that the coupling constant is nonzero $\lambda_0 \neq 0$ but infinitely small. As we have seen in Section 3, at $T = 0$ the superconducting order parameter $\kappa(\mathbf{p})$ is finite in the FC range, while

the maximum value of the superconducting gap $\Delta_1 \propto \lambda_0$ is infinitely small. Therefore, any small magnetic field $B \neq 0$ can be considered as a critical field and will destroy the coherence of $\kappa(\mathbf{p})$ and thus FC itself. To define the type of FC rearrangement, simple energy arguments are sufficient. On one hand, while the field is zero in the sample until then the state with FC is not destroyed. The energy gain ΔE_B due to removing the FC state is $\Delta E_B \propto B^2$ and tends to zero with $B \rightarrow 0$. On the other hand, occupying the finite range $(p_f - p_i)$ in the momentum space, the function $n_0(\mathbf{p})$ given by Eq. (2.8), or by Eq. (3.6), leads to a finite gain in the ground state energy [17]. Thus, a new ground state replacing FC should have almost the same energy as the former one. Such a state is given by the multiconnected Fermi spheres resembling an onion, where the smooth quasiparticle distribution function $n_0(\mathbf{p})$ in the $(p_f - p_i)$ range is replaced by a multiconnected distribution $\nu(\mathbf{p})$ [82, 83]

$$\nu(\mathbf{p}) = \sum_{k=1}^n \theta(p - p_{2k-1}) \theta(p_{2k} - p). \quad (5.1)$$

Here the parameters $p_i \leq p_1 < p_2 < \dots < p_{2n} \leq p_f$ are adjusted to obey the normalization condition:

$$\int_{p_{2k-1}}^{p_{2k+3}} \nu(\mathbf{p}) \frac{d\mathbf{p}}{(2\pi)^3} = \int_{p_{2k-1}}^{p_{2k+3}} n_0(\mathbf{p}) \frac{d\mathbf{p}}{(2\pi)^3}.$$

The corresponding multiconnected distribution is shown in Fig. 4.




FIG. 4: The function $\nu(\mathbf{p})$ for the multiconnected distribution replacing the function $n_0(\mathbf{p})$ in the range $(p_f - p_i)$ occupied by FC so that $p_i < p_F < p_f$.

For definiteness, let us consider the most interesting case of a 3D system, while the consideration of a 2D system also goes along the same line. We note that the idea of multiconnected Fermi spheres, with production of new, interior segments of the Fermi surface, has been considered some time ago [84, 85]. Let us assume that the thickness δp of each interior block is approximately the same

$\delta p \simeq p_{2k+1} - p_{2k}$ and δp is defined by B . Using the Simpson's rule, we obtain that the minimum loss in the ground state energy due to formation of the blocks is about $(\delta p)^4$. This result can be understood by considering that the continuous FC function $n_0(\mathbf{p})$ delivers the minimum value to the energy functional $E[n(\mathbf{p})]$, while the approximation of $\nu(\mathbf{p})$ by the steps of size δp produces the minimum error of the order of $(\delta p)^4$. Taking into account that the gain due to the magnetic field is proportional to B^2 and equating both of the contribution we obtain

$$\delta p \propto \sqrt{B}. \quad (5.2)$$

Thus, at $T \rightarrow 0$ when $B \rightarrow 0$, the thickness δp goes to zero as well, $\delta p \rightarrow 0$, while the NFL behavior of the FC state is replaced by the behavior of LFL with the Fermi momentum p_f . It follows from Eq. (3.6) that $p_f > p_F$, while the number density x of itinerant electrons remains constant. As we shall see, this observation plays important role when considering the Hall coefficient $R_H(B)$ as a function of B at low temperatures in the HF metals with FC.

To calculate the effective mass $M^*(B)$ as function of the applied magnetic field B , we observe that at $T = 0$ the application of finite magnetic field B splits the FC state into the Landau levels and suppresses the superconducting order parameter $\kappa(\mathbf{p})$ thus destroying the FC state. Therefore the LFL behavior is expected to be restored [8, 86]. The Landau levels at the Fermi surface can be approximated by a single block whose thickness in momentum space is δp . Approximating the dispersion of quasiparticles within this block by $\varepsilon(p) \sim (p - p_F + \delta p)(p - p_F)/M$, we obtain that the effective mass $M^*(B) \sim M/(\delta p/p_F)$. The energy loss ΔE_{FC} due to rearrangement of the FC state related to this block can be estimated using the Landau formula [27]

$$\Delta E_{FC} = \int (\varepsilon(\mathbf{p}) - \mu) \delta n(\mathbf{p}) \frac{d\mathbf{p}^3}{(2\pi)^3}. \quad (5.3)$$

The region occupied by the variation $\delta n(\mathbf{p})$ has the thickness δp , while $(\varepsilon(\mathbf{p}) - \mu) \sim (p - p_F)p_F/M^*(B)$. As a result, we have $\Delta E_{FC} \sim \delta p^2/M^*(B)$. On the other hand, there is a gain $\Delta E_B \sim (B^2\mu_B)^2 M^*(B)p_F$ due to the application of the magnetic field and coming from the Zeeman splitting. Equating ΔE_B to ΔE_{FC} and taking into account that in this case $M^*(B) \propto 1/\delta p$, we obtain the following relation

$$\frac{\delta p^2}{M^*(B)} \propto \frac{1}{(M^*(B))^3} \propto B^2 M^*(B). \quad (5.4)$$

It follows from Eq. (5.4) that the effective mass $M^*(B)$ diverges as

$$M^*(B) \propto \frac{1}{\sqrt{B - B_{c0}}}. \quad (5.5)$$

Here, B_{c0} is the critical magnetic field which drives both a HF metal to its magnetic field tuned QCP and the corresponding Néel temperature toward $T = 0$ [8]. We note that in some cases $B_{c0} = 0$, for example, the HF metal CeRu_2Si_2 shows neither evidence of the magnetic ordering, superconductivity down to the lowest temperatures nor the LFL behavior [87]. Equation (5.5) shows that by applying the magnetic field $B > B_{c0}$ the system can be driven back into LFL with the effective mass $M^*(B)$ depending on the magnetic field. This means that the following dependences are valid: for the coefficient $A(B) \propto (M^*(B))^2$ [88], for the specific heat, $C/T = \gamma_0(B) \propto M^*(B)$, and for the magnetic susceptibility $\chi_0(B) \propto M^*(B)$. The coefficient $A(B)$ determines the temperature dependent part of the resistivity, $\rho(T) = \rho_0 + \Delta\rho$, with ρ_0 being the residual resistivity and $\Delta\rho = A(B)T^2$. Since the coefficient is directly determined by the effective mass, we obtain from Eq. (5.5) that

$$A(B) \propto \frac{1}{B - B_{c0}}. \quad (5.6)$$

It is seen that the well-known empirical Kadowaki-Woods (KW) ratio [9], $K = A/\gamma_0^2 \simeq \text{const}$, is fulfilled. At this point, we stress that the value of K may be dependent on the degeneracy number of quasiparticles as it was recently shown. As a result, the grand-KW-relation produces good description of the data for whole the range of degenerate HF systems [89]. In the simplest case when the heavy electron liquid is formed by quasiparticles with spin 1/2 and the degeneracy number is 2, K turns out to be close to the empirical value [88], called as the KW relation [9]. Therefore, we come to the conclusion that by applying magnetic fields the system is driven back into the LFL state where the constancy of the Kadowaki-Woods ratio is obeyed.

At finite temperatures, the system remains in the LFL state, but there exists a temperature $T^*(B)$, at which the NFL behavior is restored. To calculate the function $T^*(B)$, we observe that the effective mass M^* characterizing the single particle spectrum cannot be changed at $T^*(B)$ since there are no any phase transitions. In other words, at the crossover point, we have to compare the effective mass $M^*(T)$ defined by $T^*(B)$, Eq. (2.12), and that $M^*(B)$ defined by the magnetic field B , Eq. (5.5), $M^*(T) \sim M^*(B)$

$$\frac{1}{M^*(T)} \propto T^*(B) \propto \frac{1}{M^*(B)} \propto \sqrt{B - B_{c0}}. \quad (5.7)$$

As a result, we obtain

$$T^*(B) \propto \sqrt{B - B_{c0}}. \quad (5.8)$$

At temperatures $T \geq T^*(B)$, the system comes back to the NFL behavior with M^* defined by Eq. (2.12), and the LFL behavior disappear. We can conclude that Eq. (5.8) determines the line in the

$B - T$ phase diagram that separates the region of the B dependent effective mass from the region of the T dependent effective mass. At the temperature $T^*(B)$, there occurs a crossover from the T^2 dependence of the resistivity to the T dependence. It follows from Eq. (5.8), that the heavy electron liquid at some temperature T can be driven back into the Landau Fermi-liquid by applying a strong enough magnetic field $(B - B_{c0}) \propto (T^*(B))^2$. We can also conclude, that at finite temperature $T < T^*(B)$, the heavy electron liquid shows a more pronounced metallic behavior at the elevated magnetic field B since the effective mass decreases (see Eq. (5.5)). The same behavior of the effective mass can be observed in the Shubnikov — de Haas oscillation measurements. We conclude that one obtains a unique possibility to control the essence of the strongly correlated liquid by magnetic fields which induce the change of the NFL behavior to the LFL liquid behavior.

Let us briefly consider the case when the system is very near FCQPT being on the ordered side and therefore $\delta p_{FC} = (p_f - p_i)/p_F \ll 1$. Since $\delta p \propto M^*(B)$, it follows from Eqs. (5.2) and (5.5) that

$$\frac{\delta p}{p_F} \sim a_c \sqrt{\frac{B - B_{c0}}{B_{c0}}}, \quad (5.9)$$

where a_c is a constant, which is expected to be of the order of a unit, $a_c \sim 1$. At bigger magnetic field B , the value of $\delta p/p_F$ becomes comparable with δp_{FC} , and the distribution function $\nu(\mathbf{p})$ vanishes being replaced by the conventional Zeeman splitting. As a result, we are dealing with the heavy electron liquid located on the disordered side of FCQPT. As we shall see in Section 9, the behavior of this system is quite different from that of the system with FC. It follows from Eq. (5.9) that relatively weak magnetic field B_{cr}

$$(\delta p_{FC})^2 \sim \frac{B_{cr} - B_{c0}}{B_{c0}}, \quad (5.10)$$

removes the system from the ordered side of the phase transition provided that $\delta p_{FC} \ll 1$.

6. MAGNETIC-FIELD INDUCED LANDAU FERMI LIQUID IN HIGH- T_c METALS

The LFL theory has revealed that the low-energy elementary excitations of a Fermi liquid look like the spectrum of an ideal Fermi gas. These excitations are described in terms of quasiparticles with an effective mass M^* , charge e and spin $1/2$. The quasiparticles define the major part of the low-temperature properties of Fermi liquids. As we have shown in Section 5, at temperatures $T < T^*(B)$, the LFL behavior of the heavy electron liquid with FC is recovered by the application of magnetic field B larger than the critical field B_c suppressing the superconductivity. Thus, the heavy electron liquid with FC can be viewed as LFL induced by the magnetic field. In such a state, the

Wiedemann-Franz (WF) law and the Korringa law are held and the elementary excitations are LFL quasiparticles. Our consideration is valid for relatively weak magnetic fields when the contributions coming from the magnetic field are proportional B^2 and the Zeeman splitting is much smaller than the Fermi momentum.

It was reported recently that in the normal state obtained by applying a magnetic field greater than the upper critical field B_c , in a hole-doped cuprates at overdoped concentration ($\text{Ti}_2\text{Ba}_2\text{CuO}_{6+\delta}$) [11] and at optimal doping concentration ($\text{Bi}_2\text{Sr}_2\text{CuO}_{6+\delta}$) [90], there are no any sizable violations of the WF law. In the electron-doped copper oxide superconductor $\text{Pr}_{0.91}\text{LaCe}_{0.09}\text{CuO}_{4-y}$ ($T_c=24$ K) when superconductivity is eliminated by a magnetic field, it was found that the spin-lattice relaxation rate $1/T_1$ follows the $T_1T = \text{constant}$ relation, known as the Korringa law [91], down to temperature of $T = 0.2$ K [92]. At higher temperatures and applied magnetic fields of 15.3 T perpendicular to the CuO_2 plane, $1/T_1T$ as a function of T is a constant below $T = 55$ K. At $300 \text{ K} > T > 50 \text{ K}$, $1/T_1T$ decreases with growing T [92]. Recent measurements for strongly overdoped non-superconducting $\text{La}_{1.7}\text{Sr}_{0.3}\text{CuO}_4$ have shown that the resistivity ρ exhibits T^2 behavior, and the WF law holds perfectly [80, 81]. Since the validity of the WF and the Korringa laws are the robust signature of LFL, these experimental facts demonstrate that the observed elementary excitations cannot be distinguished from the Landau quasiparticles. This imposes strong constraints for models describing the hole-doped and electron-doped high-temperature superconductors. For example, in the cases of a Luttinger liquid [93], spin-charge separation (see e.g. [94]), and in some solutions of $t - J$ model [95] a violation of the WF law was predicted.

As any phase transition, FCQPT is related to the order parameter, which induces a broken symmetry. It was shown in Section 3 that the relevant order parameter is the superconducting order parameter $\kappa(\mathbf{p})$, which is suppressed by the critical magnetic field B_c , when $B_c^2 \sim \Delta_1^2$. If the coupling constant $\lambda_0 \rightarrow 0$, the critical magnetic field $B_c \rightarrow 0$ will destroy the state with FC converting the strongly correlated Fermi liquid into LFL. The magnetic field plays the role of the control parameter determining the effective mass $M^*(B)$ as it follows from Eq. (5.5).

If λ_0 is finite, the critical field is also finite, and Eq. (5.5) is valid at $B > B_c$. In that case the system is driven back to LFL and has the LFL behavior induced by the magnetic field. Then, the low energy elementary excitations are characterized by $M^*(B)$ and cannot be distinguished from Landau quasiparticles. As a result, at $T \rightarrow 0$, the WF law is held in accordance with experimental facts [11, 90]. On the hand, in contrast to the LFL theory, the effective mass $M^*(B)$ depends on the magnetic field.

Equation (5.5) shows that by applying a magnetic field $B > B_c$ the system can be driven back into LFL with the effective mass $M^*(B)$ which is finite and independent of the temperature. This means that the low temperature properties depend on the effective mass in accordance with the LFL theory. In particular, the resistivity $\rho(T)$ as a function of the temperature behaves as $\rho(T) = \rho_0 + \Delta\rho(T)$ with $\Delta\rho(T) = AT^2$, and the factor $A \propto (M^*(B))^2$. Taking into account that in the case of the high- T_c superconductors B_{c0} is expected to be zero, we obtain from Eq. (5.5) that

$$\gamma_0\sqrt{B} = \text{const.} \quad (6.1)$$

Here $\gamma_0 = C/T$ with C is the specific heat. Taking into account Eqs. (5.6) and Eq. (6.1), we obtain

$$\gamma_0 \sim A(B)\sqrt{B}. \quad (6.2)$$

At finite temperatures, the system remains LFL, but there is the crossover from the LFL behavior to the non-Fermi liquid behavior at temperature $T^*(B) \propto \sqrt{B}$. At $T > T^*(B)$, the effective mass starts to depend on the temperature $M^* \propto 1/T$, and the resistivity possesses the non-Fermi liquid behavior with a substantial linear term, $\Delta\rho(T) \propto T$ [29, 30, 83]. Such a behavior of the resistivity was observed in the cuprate superconductor $\text{Ti}_2\text{Ba}_2\text{CuO}_{6+\delta}$ ($T_c < 15$ K) [96]. At $B=10$ T, $\Delta\rho(T)$ is a linear function of the temperature between 120 mK and 1.2 K, whereas at $B = 18$ T, the temperature dependence of the resistivity is consistent with $\Delta\rho(T) = AT^2$ in the same temperature range [96].

In LFL, the nuclear spin-lattice relaxation rate $1/T_1$ is determined by the quasiparticles near the Fermi level whose population is proportional to M^*T , so that $1/T_1T \propto M^*$ is a constant [91, 92]. When the superconducting state is removed by the application of a magnetic field, the underlying ground state can be seen as the field induced LFL with effective mass depending on the magnetic field. As a result, the rate $1/T_1$ follows the $T_1T = \text{constant}$ relation, that is the Korringa law is held. Unlike the behavior of LFL, as it follows from Eq. (5.5), $1/T_1T \propto M^*(B)$ decreases with increasing the magnetic field at $T < T^*(B)$. At $T > T^*(B)$, we observe that $1/T_1T$ is a decreasing function of the temperature, $1/T_1T \propto M^* \propto 1/T$. These observations are in a good agreement with the experimental facts [92]. Since $T^*(B)$ is an increasing function of the magnetic field, see Eq. (5.8), the Korringa law retains its validity to higher temperatures at elevated magnetic fields. We conclude, that at temperature $T_0 \leq T^*(B_0)$ and bigger magnetic fields $B > B_0$ the system shows a more pronounced metallic behavior, since the effective mass decreases with increasing B (see Eq. (5.5)). Such a behavior of the effective mass can be observed in the de-Haas van Alphen-Shubnikov studies, $1/T_1T$ and the resistivity measurements. These experiments can shed light on the physics of high- T_c metals and reveal relationships between high- T_c metals and heavy-electron metals [97].

The existence of FCQPT can also be revealed experimentally because at densities $x > x_{FC}$, or beyond the FCQPT point, the system should be LFL at sufficiently low temperatures [86]. Recent experimental data have shown that this liquid exists in the heavily overdoped non-superconducting compound $\text{La}_{1.7}\text{Sr}_{0.3}\text{CuO}_4$ [80, 81]. It is remarkable that up to $T = 55$ K the resistivity exhibits the T^2 behavior with no additional linear term, and the WF law is verified to within the experimental resolution [80, 81]. While at elevated temperatures, a strong deviations from the LFL behavior are observed. We anticipate that in this case the system can be again driven back to the LFL behavior by the application of sufficiently strong magnetic fields.

Thus, the mentioned above striking measurements, which were used in studies of the nature of the high- T_c superconductivity, suggest that FCQPT and the emergence of the novel quasiparticles with effective mass strongly depending on the magnetic field and temperature and resembling the Landau quasiparticles are qualities intrinsic to the electronic system of the high- T_c superconductors.

7. APPEARANCE OF FCQPT IN DIFFERENT FERMI LIQUIDS

It is widely believed that unusual properties of correlated liquids observed in the high-temperature superconductors, heavy-fermion metals, 2D ^3He and etc., are determined by quantum phase transitions. Therefore, immediate experimental studies of relevant quantum phase transitions and of their quantum critical points are of crucial importance for understanding the physics of the high- T_c superconductivity and HF systems. In case of the high- T_c superconductors, these studies are difficult to carry out, because all the corresponding area is occupied by the superconductivity. On the other hand, recent experimental data on different highly correlated Fermi liquids, when the system in question is approaching FCQPT from the disordered side, can help to illuminate both the nature of this point and the control parameter, by which this point is driven. We shall call Fermi liquids approaching FCQPT from the disordered side as highly correlated ones to distinguish them from strongly correlated liquids which have undergone FCQPT. Detailed explanations on this point are given in Section 8.

Experimental facts on high-density 2D ^3He [98, 99] show that the effective mass diverges when the density, at which 2D ^3He liquid begins to solidify, is approached [99]. Then, a sharp increase of the effective mass in a metallic 2D electron system was observed, when the density tends to the critical density of the metal-insulator transition point, which occurs at sufficiently low densities [34]. Note, that there is no ferromagnetic instability in both Fermi systems and the relevant Landau amplitude

$F_0^a > -1$ [34, 99], in accordance with the almost localized fermion model [100].

Now let us consider the divergence of the effective mass in 2D and 3D highly correlated Fermi liquids at $T = 0$, when the density x approaches FCQPT from the side of normal Landau Fermi liquid, that is from the disordered phase. First, we calculate the divergence of M^* as a function of the difference $(x - x_{FC})$ in case of 2D Fermi liquid. For this purpose we use the equation for M^* obtained in [38], where the divergence of the effective mass M^* due to the onset of FC in different Fermi liquids (such as 2D and 3D electron and ^3He liquids) was predicted. At $x \rightarrow x_{FC}$, the effective mass M^* can be approximated as

$$\frac{1}{M^*} \simeq \frac{1}{M} + \frac{1}{4\pi^2} \int_{-1}^1 \int_0^{g_0} \frac{v(q(y))}{[1 - R(q(y), \omega = 0, g)\chi_0(q(y), \omega = 0)]^2} \frac{y dy dg}{\sqrt{1 - y^2}}. \quad (7.1)$$

Here we adopt the notation $p_F \sqrt{2(1 - y)} = q(y)$ with $q(y)$ being the transferred momentum, M is the bare mass, ω is the frequency, $v(q)$ is the bare interaction, and the integral is taken over the coupling constant g from zero to its real value g_0 . In Eq. (7.1), both $\chi_0(q, \omega)$ and $R(q, \omega)$ are the linear response function of a noninteracting Fermi liquid and the effective interaction respectively. They define the linear response function of the system in question

$$\chi(q, \omega, g) = \frac{\chi_0(q, \omega)}{1 - R(q, \omega, g)\chi_0(q, \omega)}. \quad (7.2)$$

In the vicinity of the charge density wave instability, occurring at the density x_{cdw} , the singular part of the function χ^{-1} on the disordered side is of the well-known form (see e.g. [2])

$$\chi^{-1}(q, \omega, g) \simeq a(x_{cdw} - x) + b(q - q_c)^2 + c(g_0 - g), \quad (7.3)$$

where a , b and c are constants and $q_c \simeq 2p_F$ is the wavenumber of the charge density wave order. Upon substituting Eq. (7.3) into Eq. (7.1) and integrating, the equation for the effective mass M^* can be presented in the following form [31, 32]

$$\frac{1}{M^*} = \frac{1}{M} - \frac{C}{\sqrt{x - x_{cdw}}}, \quad (7.4)$$

with C being some positive constant. The behavior of the effective mass as a function of the electron number density x in a silicon MOSFET is shown in Fig. 5. The fitting parameters are $x_{cdw} = 0.7 \times 10^{-11} \text{cm}^{-2}$, $C = 2.14 \times 10^{-6} \text{cm}^{-1}$ and $x_{FC} = 0.9 \times 10^{-11} \text{cm}^{-2}$ [32]. It is seen from Fig. 5 that Eq. (7.4) describes rather good the data.

It is seen from Eq. (7.4) that M^* diverges at some point x_{FC} referred to as the critical point, at which FCQPT occurs, as a function of the difference $(x - x_{FC})$ [31, 32]

$$\frac{M^*}{M} \simeq A + \frac{B}{x - x_{FC}}, \quad (7.5)$$




FIG. 5: The ratio M^*/M in a silicon MOSFET versus the carrier number density x . The filled squares denote the Shubnikov – de Haas oscillations experimental data, the data obtained by the application of a parallel magnetic field are shown by the filled circles [34, 101].

where A and B are constants. It follows from the derivation of Eqs. (7.4) and (7.5) that their forms are independent of the bare interaction $v(q)$, which effects however A , B and x_{FC} values. This result is in agreement with Eq. (2.7) which exhibits the same type of divergence independent of the interaction. Therefore both of these equations are also applicable to 2D ^3He liquid or to another Fermi liquid. It is also seen from Eqs. (7.4) and (7.5) that FCQPT precedes the formation of charge-density waves. As a consequence of this, the effective mass diverges at high densities in case of 2D ^3He , and at low densities in case of 2D electron systems, in accordance with experimental facts [34, 99]. Note, that in both cases the difference $(x - x_{FC})$ has to be positive, because x approaches x_{FC} when the system is on the disordered side of FCQPT with the effective mass $M^*(x) > 0$. Thus, in considering the 2D ^3He liquid we have to replace $(x - x_{FC})$ by $(x_{FC} - x)$ on the right hand side of Eq. (7.5). In case of a 3D system, at $x \rightarrow x_{FC}$, the effective mass is given by [38]

$$\frac{1}{M^*} \simeq \frac{1}{M} + \frac{p_F}{4\pi^2} \int_{-1}^1 \int_0^{g_0} \frac{v(q(y))y dy dg}{[1 - R(q(y), \omega = 0, g)\chi_0(q(y), \omega = 0)]^2}. \quad (7.6)$$

The comparison of Eq. (7.1) and Eq. (7.6) shows that there is no fundamental difference between these equations, and along the same way we again arrive at Eqs. (7.4) and (7.5). The only difference between 2D electron systems and 3D ones is that in the latter FCQPT occurs at densities which are well below those corresponding to 2D systems. For bulk ^3He , FCQPT cannot probably take place




FIG. 6: The ratio M^*/M in a 2D ^3He versus the fluid number density x inferred from the heat capacity measurements and the magnetization measurements [99]. The solid line represents $M^*/M = A + \frac{B}{(x_{FC}-x)}$, with $A=1.09$, $B=1.68 \text{ nm}^{-2}$ and $x_{FC} = 5.11 \text{ nm}^{-2}$.

since it is absorbed by the first order solidification [99]. The apparent divergence of the effective mass $M^*(x)$ obtained in measurements on 2D ^3He [99] is shown in Fig. 6. This is in good agreement with the divergence given by Eqs. (7.5) and (2.7).

8. HEAVY FERMION METALS WITH HIGHLY CORRELATED ELECTRON LIQUID

In HF metals with strong electron correlations, quantum phase transitions at zero temperature may strongly influence the measurable quantities up to relatively high temperatures. These quantum phase transitions have recently attracted much attention because the behavior of HF metals is expected to follow universal patterns defined by the quantum mechanical nature of the fluctuations taking place at quantum critical points (see e.g. [3, 4]). Only recently, there appeared experimental facts which deliver experimental grounds to understand the nature of quantum phase transition producing the universal behavior of HF metals. It has been demonstrated that at low temperatures the main properties of HF metals such as the magnetoresistance, resistivity, specific heat, magnetization, susceptibility, volume thermal expansion, etc, strongly depend on temperature T and applied magnetic field B . As a result, these properties can be controlled by placing these metals at the special point of the field-temperature $B - T$ phase diagram. In the LFL theory, considered as the

main instrument when investigating quantum many electron physics, the effective mass M^* of quasiparticle excitations determining the thermodynamic properties of electronic systems is practically independent of temperature and applied magnetic fields. Therefore, the observed anomalous behavior is uncommon and can hardly be understood within the framework of the conventional LFL theory based on the notion of quasiparticles. As a result, it is necessary to use theories that are based on the Landau concept of the order parameter which is introduced to classify phases of the state of matter. These theories connect the anomalous behavior with critical fluctuations of the magnetic order parameter. These fluctuations suppressing the quasiparticles are attributed to CQPT taking place when the system in question approaches its QCP. As it was noted in Introduction, the universal behavior is only observable if the electron system of HF metal is very near QCP, for example, when the correlation length is much larger than the microscopic length scales. In the case of CQPT, the physics is dominated by thermal and quantum fluctuations of the critical state, which is characterized by the absence of quasiparticles. It is believed that the absence of quasiparticle-like excitations is the main cause of the NFL behavior and other types of the critical behavior in the quantum critical region. However, theories based on CQPT fail to explain the experimental observations related to the divergence of the effective mass M^* at the magnetic field tuned QCP, the specific behavior of the spin susceptibility and its scaling properties, the thermal expansion behavior, etc, see e.g. [6, 10, 13, 14, 87, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111].

The LFL theory rests on the notion of quasiparticles which represent elementary excitations of a Fermi liquid. Therefore these are appropriate excitations to describe the low temperature thermodynamic properties. The inability of the LFL theory to explain the experimental observations which point to the dependence of M^* upon the temperature T and applied magnetic field B may lead to the conclusion that the quasiparticles do not survive near QCP, and one might be further led to the conclusion that the heavy electron does not retain its integrity as a quasiparticle excitation (see e.g. [102, 106, 107, 112, 113]).

The mentioned above inability to explain the behavior of HF metals at QCP within the framework of theories based on CQPT may also lead to the conclusion that the other important Landau concept of the order parameter fails as well, see e.g. [106, 107, 112, 113]. Thus, we are left without the most fundamental principles of many body quantum physics while a great deal of interesting NFL phenomena related to the anomalous behavior and the experimental facts collected in measurements on the HF metals remain out of reasonable theoretical explanations.

On the other hand, it is the very nature of HF metals that suggests that their unusual properties

are defined by a quantum phase transition related to the unlimited growth of the effective mass at its QCP. Moreover, a divergence to infinity of the effective electron mass was observed at a magnetic field-induced QCP [10, 102, 105]. In Section 2, we have demonstrated that such a quantum phase transition is to be FCQPT, an essential feature of which is the divergence of the effective mass M^* at its QCP, see Eq. (2.7).

8.1. Highly correlated heavy electron liquid

When a Fermi system approaches FCQPT from the disordered phase it remains the Landau Fermi liquid with the effective mass M^* strongly depending on the distance $r = (x - x_{FC})$, temperature T and a magnetic field B . This state of the system, with M^* essentially depending on T , r and B , resembles the strongly correlated liquid described in Section 2. However, in contrast to the strongly correlated liquid, there is no energy scale E_0 given by Eq. (2.14) and the system under consideration is the Landau Fermi liquid at sufficiently low temperatures with the effective mass $M^* \propto 1/r$ (see Eqs. (2.7) and (7.5)). Therefore this liquid can be called the *highly correlated liquid* which is obviously to have uncommon properties [26, 31]. Again, we use the heavy electron liquid model to study the universal behavior of the HF metals at low temperatures. As it was mentioned in Section 2, it is possible, since we consider processes related to the power-law divergence of the effective mass. This divergence is determined by small momenta transferred as compared to momenta of the order of the reciprocal lattice cell, and the contribution coming due to the lattice structure can be ignored. Thus, we may usefully ignore the complications related to the lattice and get rid of the specific peculiarities of a HF metal regarding the medium as homogeneous heavy electron isotropic liquid.

The effective mass M^* of quasiparticle excitations controlling the density of states determines the thermodynamic properties of electronic systems. To study the behavior of the effective mass $M^*(T, B)$ as a function of temperature and magnetic field, we use the Landau equation determining $M^*(T, B)$. In the case of a homogeneous liquid, at finite temperatures and low magnetic fields, this equation reads [27]

$$\frac{1}{M^*(T, B)} = \frac{1}{M} + \sum_{\sigma_1} \int \frac{\mathbf{p}_F \mathbf{p}_1}{p_F^3} F_{\sigma, \sigma_1}(\mathbf{p}_F, \mathbf{p}_1) \frac{\partial n_{\sigma_1}(\mathbf{p}_1, T, B)}{\partial p_1} \frac{d\mathbf{p}_1}{(2\pi)^3}. \quad (8.1)$$

Here $F_{\sigma, \sigma_1}(\mathbf{p}_F, \mathbf{p}_1)$ is the Landau amplitude depending on the momenta p and spins σ , p_F is the Fermi momentum, M is the bare mass of an electron and $n_{\sigma}(\mathbf{p}, T)$ is the quasiparticle distribution function. Since HF metals are predominantly three dimensional (3D) structures we treat the homogeneous

heavy electron liquid as a 3D liquid also. For the sake of simplicity, we omit the spin dependence of the effective mass since in the case of a homogeneous liquid and weak magnetic fields, $M^*(T, B)$ does not noticeably depend on the spins. The quasiparticle distribution function is of the form

$$n_\sigma(\mathbf{p}, T) = \left\{ 1 + \exp \left[\frac{(\varepsilon(\mathbf{p}, T) - \mu_\sigma)}{T} \right] \right\}^{-1}, \quad (8.2)$$

where $\varepsilon(\mathbf{p}, T)$ is determined by Eq. (2.2). In our case, the single-particle spectrum does not noticeably depend on the spin, while the chemical potential may have a dependence due to the Zeeman splitting. We will show explicitly the spin dependence of a physical value when this dependence is of importance for understanding.

Replacing $n_\sigma(\mathbf{p}, T, B)$ by $n_\sigma(\mathbf{p}, T, B) \equiv \delta n_\sigma(\mathbf{p}, T, B) + n_\sigma(\mathbf{p}, T = 0, B = 0)$ where $\delta n_\sigma(\mathbf{p}, T, B) = n_\sigma(\mathbf{p}, T, B) - n_\sigma(\mathbf{p}, T = 0, B = 0)$, Eq. (1) takes the form

$$\frac{M}{M^*(T, B)} = \frac{M}{M^*(x)} + \frac{M}{p_F^2} \sum_{\sigma_1} \int \frac{\mathbf{p}_F \mathbf{p}_1}{p_F} F_{\sigma, \sigma_1}(\mathbf{p}_F, \mathbf{p}_1) \frac{\partial \delta n_{\sigma_1}(\mathbf{p}_1, T, B)}{\partial p_1} \frac{d\mathbf{p}_1}{(2\pi)^3}. \quad (8.3)$$

We assume that the heavy electron liquid is near FCQPT, therefore the distance r is small so that $M/M^*(x) \ll 1$, as it is seen from Eq. (2.7). In the case of normal metals with the effective mass of the order of a few bare electron masses, $M/M^*(x) \sim 1$, and up to temperatures $T \sim 100$ K, the second term on the right hand side of Eq. (8.3) is of the order of T^2/μ^2 and is much smaller than the first term. Thus, the system in question demonstrates the LFL behavior with the effective mass being practically independent of temperature, that is the corrections are proportional to T^2 . One can check that the same is true when magnetic field up to $B \sim 30$ T is applied. Near the critical point x_{FC} , when $M/M^*(x \rightarrow x_{FC}) \rightarrow 0$, the behavior of the effective mass changes drastically, because the first term on the right hand side of Eq. (8.3) vanishes and the second term determines the effective mass itself rather than small corrections to $M^*(x)$ related to T and B . In that case, Eq. (8.3) becomes a homogeneous equation and determines the effective mass as a function of B and T . As we will see, Eq. (8.3) describes both the NFL behavior and the LFL one with the presence of quasiparticles. In contrast to the conventional Landau quasiparticles these are characterized by the effective mass that strongly depends on both the magnetic field and the temperature.

Let us turn to a qualitative analysis of solutions of Eq. (8.3) when $x \simeq x_{FC}$. We start with the case when $T = 0$ and B is finite. The application of magnetic field leads to the Zeeman splitting of the Fermi surface and the difference δp between the Fermi surfaces with “spin up” and “spin down” becomes $\delta p = p_F^\uparrow - p_F^\downarrow \sim \mu_B B M^*(B)/p_F$ with μ_B being the Bohr magneton. Upon taking this into account, we observe that the second term in Eq. (8.3) is proportional to $(\delta p)^2 \propto (\mu_B B M^*(B)/p_F)^2$,

and Eq. (8.3) takes the form [8, 86, 114]

$$\frac{M}{M^*(B)} = \frac{M}{M^*(x)} + c \frac{(\mu_B B M^*(B))^2}{p_F^4}, \quad (8.4)$$

where c is a constant. Note that the effective mass $M^*(B)$ depends on x as well and this dependence disappears at $x = x_{FC}$. At the point $x = x_{FC}$, the term $M/M^*(x)$ vanishes, Eq. (8.4) becomes homogeneous and can be solved analytically [31, 86, 114]

$$M^*(B) \propto \frac{1}{(B - B_{c0})^{2/3}}. \quad (8.5)$$

Here B_{c0} is the critical magnetic field which drives both a HF metal to its magnetic field tuned QCP and the corresponding Néel temperature toward $T = 0$ [8]. We recall that in some cases $B_{c0} = 0$, for example, the HF metal CeRu_2Si_2 shows neither evidence of the magnetic ordering, superconductivity down to the lowest temperatures nor the LFL behavior [87].

Equation (8.5) shows the universal power low behavior of the effective mass which does not depend on the inter-particle interaction. We illustrate this behavior by calculations using a model functional [18, 19]

$$E[n(p)] = \int \frac{\mathbf{p}^2}{2M} \frac{d\mathbf{p}}{(2\pi)^3} + \frac{1}{2} \int V(\mathbf{p}_1 - \mathbf{p}_2) n(\mathbf{p}_1) n(\mathbf{p}_2) \frac{d\mathbf{p}_1 d\mathbf{p}_2}{(2\pi)^6}, \quad (8.6)$$

with the inter-particle interaction

$$V(\mathbf{p}) = g_0 \frac{\exp(-\beta_0 |\mathbf{p}|)}{|\mathbf{p}|}. \quad (8.7)$$

We normalized the effective mass by M , $M^* = M^*(B)/M$, temperature T_0 and magnetic field H by the Fermi energy ε_F^0 , $T = T_0/\varepsilon_F^0$, $H = (\mu_B B)/\varepsilon_F^0$, and use the dimensionless coupling constant $g = (g_0 M)/(2/p_i^2)$ and $\beta = \beta_0 p_F$. FCQPT takes place when the parameters reach their critical values, $\beta = b_c$ and $g = g_c$. On the other hand, FCQPT takes place when effective mass $M^* \rightarrow \infty$. This condition allows to relate b_c and g_c [18, 19]

$$\frac{g_c}{b_c^3} (1 + b_c) \exp(-b_c) [b_c \cosh(b_c) - \sinh(b_c)] = 1.$$

It follows from this equation that the critical point of FCQPT can be reached by changing g_0 if β_0 and p_F are fixed, or changing p_F if β_0 and g_0 are fixed, etc. For simplicity, we shall change g to reach FCQPT and investigate the properties of the system behind the critical point.

In Fig. 7 we present our calculations (triangles and squares) of the magnetic-field dependence of the effective mass at the critical point of FCQPT. At $\beta = b_c = 3$, FCQPT takes place when $g = g_c = 6.7167$. It is seen from Fig. 7 that the calculated power-low divergence of the effective mass is in accordance with Eq. (8.5).




FIG. 7: Calculated magnetic-field dependent effective mass M^* at fixed temperatures. Arrows indicate the temperatures. The vertical axis represents the normalized effective mass M^* . The horizontal axis is the normalized magnetic field H . Details of normalization are explained in the text. Solid lines represents $M^*(H) \propto H^{-2/3}$.

At densities $x > x_{FC}$, $M^*(x)$ is finite and we are dealing with the conventional Landau quasiparticles provided that the magnetic field is weak, so that $M^*(x)/M^*(B) \ll 1$ with $M^*(B)$ given by Eq. (8.5). In that case, the second term on the right hand side of Eq. (8.4) is proportional to $(BM^*(x))^2$ and represents small corrections. In the opposite case, when $M^*(x)/M^*(B) \gg 1$, the heavy electron liquid behaves as at QCP. Since in the LFL regime the main thermodynamic properties of the system is determined by the effective mass, it follows from Eq. (8.5) that we obtain a unique possibility to control the magnetoresistance, resistivity, specific heat, magnetization, volume thermal expansion, etc. At this point, we note that the large effective mass leads to the high density of states provoking a large number of states and phase transitions to emerge and compete with one another. Here we assume that these can be suppressed by the application of a magnetic field and concentrate on the thermodynamical properties.

To consider the qualitative behavior of $M^*(T)$ at elevated temperatures, we simplify Eq. (8.3) by omitting the variable B and simulating the influence of the applied magnetic field by the finite effective mass entering the denominator of the first term on the right hand side of Eq. (8.3). This effective mass becomes a function of the distance r , $M^*(r)$, which is determined by both B and $(x - x_{FC})$. If the magnetic field vanishes the distance is $r = (x - x_{FC})$. We integrate the second term

over the angle variable, then over p_1 by parts and substitute the variable p_1 by z , $z = (\varepsilon(p_1) - \mu)/T$. In the case of the flat and narrow band, we use the approximation $(\varepsilon(p_1) - \mu) \simeq p_F(p_1 - p_F)/M^*(T)$ and finally obtain

$$\frac{M}{M^*(T)} = \frac{M}{M^*(r)} + \alpha \int_0^\infty F(p_F, p_F(1 + \alpha z)) \frac{1}{1 + e^z} dz - \alpha \int_0^{1/\alpha} F(p_F, p_F(1 - \alpha z)) \frac{1}{1 + e^z} dz. \quad (8.8)$$

Here the notations are used: $F \sim Md(F^1 p^2)/dp$, the factor $\alpha = TM^*(T)/p_F^2 = TM^*(T)/(T_k M^*(r))$, $T_k = p_F^2/M^*(r)$. The Fermi momentum is defined as $\varepsilon(p_F) = \mu$. We first assume that $\alpha \ll 1$. Then omitting terms of the order of $\exp(-1/\alpha)$, we expand the upper limit of the second integral on the right hand side of Eq. (8.8) to ∞ and observe that the sum of the second and third terms represents an even function of α . These are the typical expressions with Fermi-Dirac functions as integrands. They can be calculated using standard procedures (see e.g. [115]). Since we need only an estimation of the integrals, we represent Eq. (8.8) as

$$\frac{M}{M^*(T)} \simeq \frac{M}{M^*(r)} + a_1 \left(\frac{TM^*(T)}{T_k M^*(r)} \right)^2 + a_2 \left(\frac{TM^*(T)}{T_k M^*(r)} \right)^4 + \dots \quad (8.9)$$

Here a_1 and a_2 are constants of the order of units. Equation (8.9) can be considered as a typical equation of the LFL theory with the only exception being the effective mass $M^*(r)$ which strongly depends on the distance $r = x - x_{FC} \geq 0$ and diverges at $r \rightarrow 0$. Nonetheless, it follows from Eq. (8.9) that when $T \rightarrow 0$ the corrections to $M^*(r)$ start with the T^2 terms provided that

$$\frac{M}{M^*(r)} \gg \left(\frac{TM^*(T)}{T_k M^*(r)} \right)^2 \simeq \frac{T^2}{T_k^2}, \quad (8.10)$$

and the system exhibits the LFL behavior. It is seen from Eq. (10) that when $r \rightarrow 0$, $M^*(r) \rightarrow \infty$, and the LFL behavior disappears. The free term on the right hand side of Eq. (8.8) vanishes, $M/M^*(r) \rightarrow 0$, and Eq. (8.8) in itself becoming homogeneous determines the value and universal behavior of the effective mass.

At some temperature $T_1 \ll T_k$, the value of the sum on the right hand side of Eq. (8.9) is determined by the second term. Then Eq. (8.10) is not valid, and upon keeping only the second term in Eq. (8.9) this can be used to determine $M^*(T)$ in a transition region [114, 116]

$$M^*(T) \propto \frac{1}{T^{2/3}}. \quad (8.11)$$

The variation as $T^{-2/3}$ exponent with the temperature growth deserves a comment. Equation (8.11) is valid if the second term in Eq. (8.9) is much larger than the first one, that is

$$\frac{T^2}{T_k^2} \gg \frac{M}{M^*(r)}, \quad (8.12)$$

and this term is grater than the third one,

$$\frac{T}{T_k} \ll \frac{M^*(r)}{M^*(T)} \simeq 1. \quad (8.13)$$

Obviously, both Eqs. (8.12) and (8.13) can be simultaneously satisfied if $M/M^*(r) \ll 1$ and T is finite. The range of temperatures, over which Eq. (8.11) is valid, shrinks to zero as soon as $r \rightarrow 0$ because $T_k \rightarrow 0$. Thus, if the system is very near QCP, $x \rightarrow x_{FC}$, it is possible to observe the behavior of the effective mass given by Eq. (8.11) in a wide range of temperatures provided that the effective mass $M^*(r)$ is diminished by the application of high magnetic fields, that is, the distance r becomes larger under the action of B . When r is finite the $T^{-2/3}$ behavior can be observed at relatively high temperatures. To estimate the transition temperature $T_1(B)$, we observe that the effective mass is a continuous function of the temperature, thus $M^*(B) \sim M^*(T_1)$. Taking into account Eqs. (8.7) and (8.11), we obtain $T_1(B) \propto B$.




FIG. 8: Calculated entropy $S(T)$ as a function of temperature at fixed magnetic field $H = 0.01$. Solid line represents $S(T)$ at the transition region $T = T_1(B)$ (left panel). At low temperatures $T < 0.007$, the system exhibits the LFL behavior, $S(T) \propto T$. The other solid (right panel) line represents $S(T) \propto T^{1/2}$.

Then, at elevated temperatures, the system enters into a different regime. The coefficient α becomes $\alpha \sim 1$, the upper limit of the second integral in Eq. (8.8) cannot be expanded to ∞ , and odd terms come into play. As a result, Eq. (8.9) is no longer valid, but the sum of both the first integral and the second one on the right hand side of Eq. (8.8) is proportional to $M^*(T)T$. Upon omitting the first term $M/M^*(r)$ and approximating the sum of the integrals by $M^*(T)T$, we solve

Eq. (8.8) and obtain

$$M^*(T) \propto \frac{1}{\sqrt{T}}. \quad (8.14)$$

We illustrate the above consideration by numerical calculations (shown by filled squares in Fig. 8) based on the model functional (8.6). In Fig. 8, we show the evolution of the low temperature entropy from the transition region $T \sim T_1(B)$ to $S(T) \propto \sqrt{T}$ at higher temperatures upon applying magnetic field $H = 0.01$. The calculated behavior of $S(T)/T \propto M^*(T)$ is in accord with Eq. (8.14). Details of the normalization T and H are given at Fig. 7.

Thus, we can conclude that at higher temperatures when $x \simeq x_{FC}$ the system exhibits three types of regimes: the LFL behavior at $\alpha \ll 1$, when Eq. (8.10) is valid; the $M^*(T) \propto T^{-2/3}$ behavior and $S(T) \propto M^*(T)T \propto T^{1/3}$, when Eqs. (8.12) and (8.13) are valid; and the $1/\sqrt{T}$ behavior of the effective mass at $\alpha \sim 1$, while the entropy $S(T) \propto M^*(T)T \propto \sqrt{T}$, and the specific heat $C(T) = T(\partial S(T)/\partial T) \propto \sqrt{T}$.

In the absence of magnetic field, calculated evolutions of $M^*(T)$, $S(T)$ and $C(T)$ based on the model functional (8.6) are shown by filled squares in Fig. 9, Fig. 10, and Fig. 11 respectively.




FIG. 9: Calculated effective mass $M^*(T)$ as a function of temperature. Solid line represents $M^*(T) \propto 1/\sqrt{T}$.

Let us estimate the quasiparticles width $\gamma(T)$. Within the framework of the LFL theory it is given by [27]

$$\gamma \sim |\Gamma|^2 (M^*)^3 T^2, \quad (8.15)$$

where Γ is the particle-hole amplitude. In the case of a strongly correlated system with its large density of states related to the huge value of the effective mass, the amplitude Γ cannot be approxi-




FIG. 10: Calculated entropy $S(T)$ as a function of temperature. Solid line represents $S(T) \propto \sqrt{T}$.




FIG. 11: Calculated specific heat $C(T)$ as a function of temperature. Solid line represents $C(T) \propto \sqrt{T}$.

mated by the bare particle interaction but can be estimated within the ladder approximation which gives $|\Gamma| \sim 1/(p_F M^*(T))$ [20]. As a result, we obtain that in the LFL regime $\gamma(T) \propto T^2$, in the $T^{-2/3}$ regime $\gamma(T) \propto T^{4/3}$, and in the $1/\sqrt{T}$ regime $\gamma(T) \propto T^{3/2}$. We observe that in all the cases the width is small as compared to the quasiparticle characteristic energy which is of the order of T , so the notion of a quasiparticle is meaningful. We can conclude that when the heavy electron liquid is near the QCP of FCQPT, being on the disordered side, its low energy excitations are quasiparticle excitations with the effective mass $M^*(B, T)$. At this point we note that at $x \rightarrow x_{FC}$, the quasiparticle renormalization factor $a(\mathbf{p})$ remains finite and approximately constant, and the divergence of the

effective mass given by Eq. (2.7) is not related to vanishing $a(\mathbf{p})$ [117]. Thus the notion of the quasi-particles is preserved and these are the relevant excitations when considering the thermodynamical properties of the heavy electron liquid.

8.2. Resistivity of heavy fermion metals

Since the resistivity, $\rho(T) = \rho_0 + \Delta\rho(T)$, is directly determined by the effective mass, because the coefficient $A(B, T) \propto (M^*(B, T))^2$ [88], the above mentioned temperature dependences can be observed in measurements of the resistivity of HF metals.

At $T \ll T_1$, the system in question demonstrates the LFL regime, the divergence of the effective mass at $x \rightarrow x_{FC}$ is described by Eq. (8.5) and the coefficient $A(B)$ diverges as

$$A(B) \propto \frac{1}{(B - B_{c0})^{4/3}}. \quad (8.16)$$

Thus, the resistivity, $\rho(B) \propto A(B)T^2$, as a function of magnetic field diverges as given by Eq. (8.16). As a function of temperature, the resistivity behaves as $\Delta\rho_1 = c_1 T^2 / (B - B_{c0})^{4/3} \propto T^2$. The second is NFL regime which is determined by Eq. (8.11) and characterized by $\Delta\rho_2 = c_2 T^2 / (T^{2/3})^2 \propto T^{2/3}$. At $T > T_1(B)$, the third NFL regime is given by Eq. (8.14) and represented by $\Delta\rho_3 = c_3 T^2 / (\sqrt{T})^2 \propto T$. Here c_1, c_2, c_3 are constants. It is remarkable that all temperature dependences corresponding to these regimes were observed in measurements on the HF metals CeCoIn₅ and YbAgGe [14, 104, 105, 109]. If we consider the ratio $\Delta\rho_2/\Delta\rho_1 \propto ((B - B_{c0})/T)^{4/3}$, we come to a very interesting conclusion that the ratio is a function of only the variable $(B - B_{c0})/T$, thus representing the scaling behavior. This result is in excellent agreement with experimental facts [105].

8.3. Magnetic susceptibility

The magnetic susceptibility is proportional to the effective mass, $\chi \propto M^*$, with M^* given by Eq. (8.5). Therefore, at $T \ll T_1$,

$$\chi(B) \propto M^*(B) \propto (B - B_{c0})^{-2/3}, \quad (8.17)$$

while the static magnetization $M_B(B)$ is given by

$$M_B(B) \propto B M^*(B) \propto (B - B_{c0})^{1/3}. \quad (8.18)$$

At $T \gg T_1$, as it follows from Eq. (8.14), Eq. (8.17) has to be rewritten as

$$\chi(T) \propto M^*(T) \propto \frac{1}{\sqrt{T}}. \quad (8.19)$$

The observed behavior of $\chi(B)$ and $M_B(B)$ and the behavior of $\chi(T)$ are in accord with results of measurements on CeRu₂Si₂ with the critical field $B_{c0} \rightarrow 0$ [87].

Consider the state of the system when $r \rightarrow 0$. Its properties are determined by magnetic fields B and temperature T because there are no other parameters to describe such a state. At the transition temperatures $T \simeq T_1(B)$, the effective mass depends on both T and B , while at $T \ll T_1(B)$, the system is LFL with the effective mass being given by Eq. (8.5), and at $T \geq T_1(B)$, the mass is defined by Eq. (8.11). Instead of solving Eq. (8.3), it is possible to construct a simple interpolation formula to describe the behavior of the effective mass over all regions,

$$M^*(B, T) = \frac{1}{c_1(B - B_{c0})^{2/3} + c_2 f(y) T^{2/3}}. \quad (8.20)$$

Here, $f(y)$ is a universal monotonic function of $y = (T/(B - B_{c0}))^{2/3}$ such that $f(y \sim 1) = 1$, and $f(y \ll 1) = 0$. It is seen from Eq. (8.20) that the behavior of the effective mass can be represented by a universal function of only one variable y if the temperature is measured in the units of the transition temperature $T_1(B)$, and the effective mass is measured in the units of $M^*(B)$ given by Eq. (8.5). This representation describes the scaling behavior of the effective mass. As seen from Eqs. (8.18) and (8.20), the scaling behavior of the magnetization can be represented in the same way, provided the magnetization is normalized by the saturated value at each field given by Eq. (8.5)

$$\frac{M_B(B, T)}{M_B(B)} \propto \frac{1}{1 + c_3 f(y) y}, \quad (8.21)$$

where c_3 is a constant. It is seen from Eq. (8.21), that magnetization is a monotonic function of y . Upon using the definition of susceptibility, $\chi = \partial M_B / \partial B$, we come to the conclusion that the susceptibility also exhibits the scaling behavior and can be presented as a universal function of only one variable y , provided it is normalized by the saturated value at each field given by Eq. (8.17)

$$\frac{\chi(B, T)}{\chi(B)} \propto \frac{1}{1 + c_3 f(y) y} + 2c_3 y \frac{f(y) + y df(y)/dy}{(1 + c_3 f(y) y)^2}. \quad (8.22)$$

It is of importance to note that the susceptibility is not a monotonic function of y because the derivative is the sum of two contributions with different behavior. The second contribution on the right hand side of Eq. (8.22) gives the susceptibility a maximum [8, 114, 116]. As shown in Fig. 12, the above behaviors of the magnetization and susceptibility are in accord with numerical calculations of the susceptibility and magnetization [114] and with the facts observed in measurements on CeRu₂Si₂ [87].

It is seen from Fig. 12 that for finite B , the curve describing $\chi(B, T)$ acquires a maximum at some temperature T_P . This behavior is in good agreement with experimental facts [87] and associated with




FIG. 12: Normalized magnetic susceptibility $\chi(B, T)/\chi(B, T_P)$ (left panel) and normalized magnetization $M_B(B, T)/M_B(B, T_P)$ (right panel) for CeRu_2Si_2 in magnetic fields 0.20 mT (squares), 0.39 mT (triangles), and 0.94 mT (circles), plotted against normalized temperature T/T_P [87], where T_P is the temperature at peak susceptibility. The solid curves trace the calculated universal behavior [114].

the suppression of the divergent NFL terms $\sim T^{-2/3}$ in $\chi(B, T)$ and recovery of the LFL behavior at static magnetic fields in which the Zeeman energy splitting $\mu_B B$ exceeds T [114]. Note that magnetization $M_B(B, T)$ does not exhibit a maximum. In Fig. 12, the temperature is normalized by T_P , the susceptibility is normalized by its peak height $\chi(B, T_P)$ and the magnetization by the saturated value at each field.

8.4. Magnetoresistance

Using the results just presented, we consider the behavior of magnetoresistance (MR)

$$\rho_{mr}(B, T) = \frac{\rho(B, T) - \rho(0, T)}{\rho(0, T)}, \quad (8.23)$$

as a function of magnetic field B and T . Here the resistivity, $\rho(B, T) = \rho_0 + \Delta\rho(B, T) + \Delta\rho_{mr}(B)$, is measured at the magnetic field B and temperature T . We assume that the contribution $\Delta\rho_{mr}(B)$ coming from the magnetic field B can be treated within the low field approximation and given by the well-known Kohler's rule,

$$\frac{\Delta\rho_{mr}(B)}{\rho(0, T)} \simeq \lambda_{\perp} \left(\frac{B\rho(0, \Theta_D)}{B_0\rho(0, T)} \right)^2, \quad (8.24)$$

with Θ_D is the Debye temperature, B_0 is the characteristic field and $\lambda_{\perp}$ is a constant. Note, that the low field approximation implies that $\Delta\rho_{mr}(B) \ll \rho(0, T) \equiv \rho(T)$. We also assume that that

temperature is not too low so that $\rho_0 \leq \Delta\rho(B=0, T)$, while $B \geq B_{c0}$. Substituting Eq. (8.24) into Eq. (8.23), we find that

$$\rho_{mr}(B, T) \sim \frac{c(M^*(B, T))^2 T^2 + \Delta\rho_{mr}(B) - c(M^*(0, T))^2 T^2}{\rho(0, T)}. \quad (8.25)$$

Here $M^*(B, T)$ denotes the effective mass which now depends on both the magnetic field and the temperature, and c is a constant determining the temperature dependent part of the resistivity, $c(M^*(B, T))^2 T^2 = \Delta\rho(B, T)$.

Consider MR given by Eq. (8.25) as a function of B at some temperature $T = T_0$. At low magnetic fields when $T_0 > T_1(B) \propto B$, the main contribution to MR comes from $\Delta\rho_{mr}(B)$ since the effective mass depends mainly on temperature. Therefore, the ratio $|M^*(B, T) - M^*(0, T)|/M^*(0, T) \ll 1$, the main contribution is given by $\Delta\rho_{mr}(B)$, and MR is an increasing function of B . When B becomes so large that $T_1(B) \sim T_0$, the difference $(M^*(B, T) - M^*(0, T))$ becomes negative, and MR as the function of B reaches its maximum value at $T_1(B) \sim T_0$. We recall that $T_1(B)$ determines the crossover from T^2 dependence of the resistivity to the T dependence. At elevated B when $T_1(B) > T_0$, the ratio $(M^*(B, T) - M^*(0, T))/M^*(0, T) \sim -1$ and MR becomes negative being a decreasing function of B .

Consider now MR as a function of T at some B_0 . At $T \leq T_1(B_0)$, we have LFL. At low temperatures $T \ll T_1(B_0)$, it follows from Eqs. (8.5) and (8.14) that $M^*(B_0)/M^*(T) \ll 1$, and MR is determined by the resistivity $\rho(0, T)$. Note, that B_0 has to be comparatively high to ensure the inequality, $M^*(B_0)/M^*(T) \ll 1$. As a result, $\rho_{mr}(B_0, T \rightarrow 0) \sim -1$, because $\Delta\rho_{mr}(B)/\rho(0, T) \ll 1$. Differentiating the function $\rho_{mr}(B_0, T)$ with respect to B_0 we can check that its slope becomes steeper as B_0 is decreased, being proportional $\propto (B_0 - B_{c0})^{-7/3}$. At $T \simeq T_1(B_0)$, MR possesses a node because at this point the effective mass $M^*(B_0) \simeq M^*(T)$, and $\rho(B_0, T) \simeq \rho(0, T)$. We can conclude that the crossover from the T^2 resistivity to the T resistivity, which occurs at $T \sim T_1(B_0)$, manifests itself in the transition from negative MR to positive MR. At $T \geq T_1(B_0)$, the main contribution to MR comes from $\Delta\rho_{mr}(B_0)$, and MR reaches its maximum value. Upon using Eqs. (8.14) and (8.25) and taking into account that at this point $T \propto (B_0 - B_{c0})$, we obtain that the maximum value $\rho_{mr}^m(B_0)$ of MR is $\rho_{mr}^m(B_0) \propto 1/(B_0 - B_{c0})$. Thus, the maximum value is a decreasing function of B_0 . At $T_1(B_0) \ll T$, MR is a decreasing function of the temperature. At these temperatures, MR becomes small comparatively to its maximum value $\rho_{mr}^m(B_0)$ because $|M^*(B, T) - M^*(0, T)|/M^*(0, T) \ll 1$ and $\Delta\rho_{mr}(B_0)/\rho(T) \ll 1$.

The recent paper [105] reports the measurements of the CeCoIn₅ resistivity in magnetic fields.

The both transitions from negative MR to positive MR with increasing T and from positive MR to negative one with increasing B were observed [105]. Thus, the above observed behavior of MR is in good agreement with the experimental facts. We believe that an additional analysis of the data [105] can reveal that the crossover from T^2 dependence of the resistivity to the T dependence occurs at $T \propto (B - B_{c0}) \div (B - B_{c0})^{4/3}$, this analysis could reveal the above described supplementary peculiarities of MR as well [26].

9. HF METALS WITH STRONGLY CORRELATED ELECTRONIC LIQUID

As we have seen in Section 2, at $T = 0$ when $r = (x - x_{FC}) \rightarrow 0$, the effective mass $M^*(r) \rightarrow \infty$ and eventually beyond the critical point $x = x_{FC}$ the distance r becomes negative making the effective mass negative as follows from Eq. (2.7). As it was shown in Section 2, the system is to undergo FCQPT. Therefore behind the critical point x_{FC} of this transition, the quasiparticle distribution function represented by the step function does not deliver the minimum to the Landau functional $E[n(\mathbf{p})]$. As a result, at $x < x_{FC}$ the quasiparticle distribution is determined by Eq. (2.8) to search the minimum of a functional, which determines the quasiparticle distribution function $n_0(\mathbf{p})$ delivering the lowest possible value to the ground state energy E . It was shown in Section 3 that the relevant order parameter $\kappa(\mathbf{p}) = \sqrt{n_0(\mathbf{p})(1 - n_0(\mathbf{p}))}$ is at the same time the order parameter of the superconducting state when the maximum value Δ_1 of the superconducting gap is infinitely small. Thus, this state cannot exist at any finite temperatures and driven by the parameter x : at $x > x_{FC}$ the system is on the disordered side of FCQPT, while at $x < x_{FC}$, the system is on the ordered side. In Section 2 we have shown that this ordered state has a strong impact on the properties of the system at finite temperatures $T \ll T_f$: the effective mass $M^*(T)$ diverges as $1/T$, see Eq. (2.12), and the electronic system with FC is characterized by the energy scale E_0 given by Eq. (2.14). As a result, we can consider FCQPT as the phase transition, which separates the *highly correlated* heavy electron liquid from the *strongly correlated* one. It was shown in Section 8, that the highly correlated liquid at $T \rightarrow 0$ and $x > x_{FC}$ behaves as LFL, therefore, FCQPT separates the regions of LFL and strongly correlated liquids. Obviously, the strongly correlated electron liquid demonstrates the NFL behavior down to zero temperature.

At $0 < T \ll T_f$, the function $n_0(\mathbf{p})$ determines the entropy $S_{NFL}(T)$ of the heavy electron liquid in its NFL state. Inserting into Eq. (2.3) the function $n_0(\mathbf{p})$, one can check that behind the point of FCQPT there is a temperature independent contribution $S_0(r) \sim (p_f - p_i)/p_F \sim |r|/x_{FC}$,

where $r = x - x_{FC}$. Another specific contribution is related to the spectrum $\varepsilon(\mathbf{p})$ which insures the connection between the dispersionless region ($p_f - p_i$) occupied by FC and the normal quasiparticles located at $p < p_i$ and at $p > p_f$, and therefore it is of the form $\varepsilon(\mathbf{p}) \propto (p - p_f)^2 \sim (p_i - p)^2$. Such a form of the spectrum can be verified in exactly solvable models for systems with FC and leads to the contribution of this spectrum to the specific heat $C \sim \sqrt{T/T_f}$ [17]. Thus at $0 < T \ll T_f$, the entropy can be approximated as

$$S_{NFL}(T) \simeq S_0(r) + a\sqrt{\frac{T}{T_f}} + b\frac{T}{T_f}, \quad (9.1)$$

with a and b are constants. The third term on the right hand side of Eq. (9.1) comes from the contribution of the temperature independent part of the spectrum $\varepsilon(\mathbf{p})$ and gives a relatively small contribution to the entropy.

The calculated evolution of $S(T)$, $C(T)$ and $M^*(T)$ based on the model functional (8.6) are shown by filled symbols in Fig. 13, Fig. 14, and Fig. 15 respectively. The calculations were carried out for $g = 7, 8, 12$ and $\beta = b_c = 3$, while the critical value $g = g_c = 6.7167$.




FIG. 13: Calculated entropy $S(T)$ as a function of temperature. Solid lines represent $S(T)$ given by Eq. (9.1).

It is seen from Fig. 13 that $S_0(r)$ increases when the system being on the ordered side moves away from FCQPT.

Obviously, the term $S_0(r)$ does not contribute to the specific heat. As a result the specific heat demonstrates the anomalous behavior $C(T) \propto \sqrt{T}$ as it is seen from Fig. 14. As to the effective




FIG. 14: Calculated specific heat $C(T)$ as a function of temperature. Solid lines represent $C(T) \propto \sqrt{T}$.




FIG. 15: Calculated effective mass $M^*(T)$ as a function of temperature. Solid lines represent $M^*(T) \propto a_1/T + a_2\sqrt{T} + a_3$.

mass $M^*(T) \propto S(T)/T$, it demonstrates the divergence $M^*(T) \propto 1/T$ in accordance with Eq. (2.5) and contains all terms defining the behavior of the entropy, as shown in Fig. 15.

9.1. The Grüneisen ratio and Hall coefficient in heavy fermion metals; $T-B$ phase diagram

The temperature independent term $S_0(r)$ determines the specific NFL behavior of the system. For example, the existence of the temperature independent term $S_0(r)$ of the entropy can be illuminated

by calculating the thermal expansion coefficient $\alpha(T)$ [7, 118], which is given by [27]

$$\alpha(T) = \frac{1}{3} \left(\frac{\partial(\log V)}{\partial T} \right)_P = -\frac{x}{3K} \left(\frac{\partial(S/x)}{\partial x} \right)_T. \quad (9.2)$$

Here, P is the pressure and V is the volume. The compressibility K is not expected to be singular at FCQPT and in systems with FC, because FC is attached to the Fermi level, and it moves along as $\mu(x)$ changes, while the compressibility $K = d\mu/d(Vx)$ is approximately constant [119]. Inserting Eq. (9.1) into Eq. (9.2), we find that

$$\alpha_{FC}(T) \simeq a_0 \sim \frac{M_{FC}^* T}{p_F^2 K}. \quad (9.3)$$

Here, a_0 is a number independent of temperature. When deriving Eq. (9.3) we keep only the main contribution coming from $S_0(r)$. On the other hand, the specific heat

$$C(T) = T \frac{\partial S(T)}{\partial T} \simeq \frac{a}{2} \sqrt{\frac{T}{T_f}}. \quad (9.4)$$

As a result, the Grüneisen ratio $\Gamma(T)$ diverges as

$$\Gamma(T) = \frac{\alpha(T)}{C(T)} \simeq 2 \frac{a_0}{a} \sqrt{\frac{T_f}{T}}. \quad (9.5)$$

At this point, we consider how the behavior of the effective mass given by Eqs. (2.12) and (8.14) correspond to experimental observations. It was recently observed that the thermal expansion coefficient $\alpha(T)/T$ measured on CeNi_2Ge_2 shows a $1/\sqrt{T}$ divergence over two orders of magnitude in the temperature range from 6 K down to at least 50 mK, while measurements on $\text{YbRh}_2(\text{Si}_{0.95}\text{Ge}_{0.05})_2$ demonstrate that $\alpha/T \propto 1/T$ [6], contrary to the LFL theory which yields $\alpha(T)/T \propto M^* \simeq \text{const.}$ Since the effective mass depends on T , we obtain that the $1/\sqrt{T}$ behavior, Eq. (8.14), is in excellent agreement with the result for the former system [7], and the $1/T$ behavior, Eq. (2.12), predicted in [118] corresponds to the latter HF metal.

We see that at $0 < T \ll T_f$, the heavy electron liquid with FC behaves as if it were placed at QCP. In fact it is placed at the quantum critical line $x < x_{FC}$, that is the critical behavior is observed at $T \rightarrow 0$ for all $x \leq x_{FC}$. At $T \rightarrow 0$, the heavy electron liquid undergoes a first-order quantum phase transition because the entropy is not a continuous function: at finite temperatures the entropy is given by Eq. (9.1), while $S(T = 0) = 0$. Therefore, the entropy undergoes a sudden jump $\delta S = S_0(r)$ in the zero temperature limit. We make up a conclusion that due to the first order phase transition, the critical fluctuations are suppressed at the quantum critical line and the corresponding divergences, for example the divergence of $\Gamma(T)$, are determined by the quasiparticles

rather than by the critical fluctuations as one could expect in the case of CQPT, see e.g. [4]. Note that according to the well known inequality, $\delta Q \leq T\delta S$, the heat δQ of the transition from the ordered phase to the disordered one is equal to zero, because $\delta Q \leq S_0(r)T \rightarrow 0$ at $T \rightarrow 0$.

To study the $B-T$ phase diagram of the heavy electron liquid with FC, we consider the case when the NFL behavior arises by the suppression of the antiferromagnetic (AF) phase upon applying a magnetic field B , for example, as it takes place in the HF metals YbRh_2Si_2 and $\text{YbRh}_2(\text{Si}_{0.95}\text{Ge}_{0.05})_2$ [6, 10]. The AF phase is represented by the heavy electron LFL, with the entropy vanishing as $T \rightarrow 0$. For magnetic fields exceeding the critical value B_{c0} at which the Néel temperature $T_N(B \rightarrow B_{c0}) \rightarrow 0$ the weakly ordered AF phase transforms into weakly polarized heavy electron LFL. As it was discussed in Section 5, at $T = 0$ the application of the magnetic field B splits the FC state occupying the region $(p_f - p_i)$ into the Landau levels and suppresses the superconducting order parameter $\kappa(\mathbf{p})$ destroying the FC state. Such a state is given by the multiconnected Fermi sphere, where the smooth quasiparticle distribution function $n_0(\mathbf{p})$ in the $(p_f - p_i)$ range is replaced by a multiconnected distribution $\nu(\mathbf{p})$, see Fig. 4. Therefore the LFL behavior is restored being represented by the weakly polarized heavy electron LFL and characterized by quasiparticles with the effective mass $M^*(B)$ given by Eq. (5.5). At elevated temperatures $T > T^*(B - B_{c0}) \propto \sqrt{B - B_{c0}}$, the NFL state is restored and the entropy of the heavy electron liquid is given by Eq. (9.1). This behavior is displayed in the $T - B$ phase diagram shown in Fig. 16.

In accordance with experimental facts we assume that at relatively high temperatures $T/T_{NO} \sim 1$ the AF phase transition is of the second order [10], where T_{NO} is the Néel temperature in the absence of the magnetic field. In that case, the entropy and the other thermodynamic functions are continuous functions at the transition temperature $T_N(B)$. This means that the entropy of the AF phase $S_{AF}(T)$ coincides with the entropy of the NFL state given by Eq. (9.1),

$$S_{AF}(T \rightarrow T_N(B)) = S_{NFL}(T \rightarrow T_N(B)). \quad (9.6)$$

Since the AF phase demonstrates the LFL behavior, that is $S_{AF}(T \rightarrow 0) \rightarrow 0$, Eq. (10) cannot be satisfied at sufficiently low temperatures $T \leq T_{crit}$ due to the temperature-independent term $S_0(r)$, see Eq. (9.1). Thus, the second order AF phase transition becomes the first order one at $T = T_{crit}$ as it is shown in Fig. 16. At $T = 0$, the critical field B_{c0} , at which the AF phase becomes the heavy LFL, is determined by the condition that the ground state energy of the AF phase coincides with the ground state energy $E[n_0(\mathbf{p})]$ of the heavy LFL, that is the ground state of the AF phase becomes degenerated at $B = B_{c0}$. Therefore, the Néel temperature $T_N(B \rightarrow B_{c0}) \rightarrow 0$, and the behavior of




FIG. 16: $T - B$ phase diagram of the heavy electron liquid. The $T_N(B/B_{c0})$ curve represents the field dependence of the Néel temperature. Line separating the antiferromagnetic (AF) and the non-Fermi liquid (NFL) state is a guide to the eye. The black dot at $T = T_{crit}$ marked by the arrow is the critical temperature, at which the second order AF phase transition becomes the first one. At $T < T_{crit}$, the thick solid line represents the field dependence of the Néel temperature when the AF phase transition is of the first order. The NFL state is characterized by the entropy S_{NFL} given by Eq. (9.1). The line separating the NFL state and the weakly polarized heavy electron Landau Fermi Liquid (LFL) is $T^*(B/B_{c0} - 1) \propto \sqrt{B/B_{c0} - 1}$.

the effective mass $M^*(B \geq B_{c0})$ is given by Eq. (5.5), that is $M^*(B)$ diverges when $B \rightarrow B_{c0}$.

We note that the corresponding quantum and thermal critical fluctuations vanish at $T < T_{crit}$ because we are dealing with the first order AF phase transition. We can also reliably conclude that the critical behavior observed at $T \rightarrow 0$ and $B \rightarrow B_{c0}$ is determined by the corresponding quasiparticles rather than by the critical fluctuations accompanying second order phase transitions. When $r \rightarrow 0$ the heavy electron liquid approaches FCQPT from the ordered phase. Obviously, $T_{crit} \rightarrow 0$ at the point $r = 0$, and we are led to the conclusion that the Néel temperature vanishes at the point when the AF second order phase transition becomes the first order one. As a result, one can expect that the contributions coming from the corresponding critical fluctuations can only lead to the logarithmic corrections to the Landau theory of the phase transitions [115], and the power low critical behavior is again defined by the corresponding quasiparticles. Thus, we conclude that the Landau paradigm based on the notions of the quasiparticles and order parameter is applicable when considering the heavy electron liquid.

Now we are in position to consider the recently observed jump in the Hall coefficient at $B \rightarrow B_{c0}$

in the zero temperature limit [120]. At $T = 0$, the application of the critical magnetic field B_{c0} suppressing the AF phase (with the Fermi momentum $p_{AF} \simeq p_F$) restores the LFL with the Fermi momentum $p_f > p_F$. At $B < B_{c0}$, the ground state energy of the AF phase is lower than that of the heavy LFL, while at $B > B_{c0}$, we are dealing with the opposite case, and the heavy LFL wins the competition. At $B = B_{c0}$, both AF and LFL have the same ground state energy being degenerated. Thus, at $T = 0$ and $B = B_{c0}$, the infinitesimal change in the magnetic field B leads to the finite jump in the Fermi momentum because the distribution function becomes multiconnected, see Fig. 4, while the overall Fermi volume remains constant. That is, the number of itinerant electrons does not change. In response the Hall coefficient $R_H(B) \propto 1/x \propto 1/p_f^3$ undergoes the corresponding sudden jump. Here we have assumed that the low temperature $R_H(B)$ can be considered as a measure of the Fermi volume and, therefore, as a measure of the Fermi momentum [120]. As a result, we obtain

$$\frac{R_H(B = B_{c0} - \delta)}{R_H(B = B_{c0} + \delta)} \simeq 1 + 3 \frac{p_f - p_F}{p_F} \simeq 1 + d \frac{S_0(r)}{x_{FC}}. \quad (9.7)$$

Here δ is an infinitesimal magnetic field, $S_0(r)/x_{FC}$ is the entropy per one heavy electron, and d is a constant, $d \sim 5$. It follows from Eq. (9.1) that the abrupt change in the Hall coefficient tends to zero when $r \rightarrow 0$ and vanishes when the system in question is on the disordered side of FCQPT [121].

Now consider the magnetic susceptibility which is proportional to the effective mass given by Eq. (5.5). Therefore, at $T \ll T^*(B)$, the magnetic susceptibility given by Eq. (5.8) is of the form

$$\chi(B) \propto M^*(B) \propto \frac{1}{\sqrt{B - B_{c0}}}, \quad (9.8)$$

while the static magnetization $M(B)$ is given by [8]

$$M(B) \propto \sqrt{B - B_{c0}}. \quad (9.9)$$

As seen from Fig. 17, the field dependence of $M(B)$ given by Eq. (9.9) is in good agreement with the data obtained in measurements on $\text{YbRh}_2(\text{Si}_{0.95}\text{Ge}_{0.05})_2$ [102]. We can also conclude that Eqs. (8.21) and (8.22) determining the scaling behavior of the effective mass, static magnetization and the susceptibility are also valid in the case of strongly correlated liquid, but the variable y is now given by $y = T/\sqrt{B - B_{c0}}$, while the function $f(y)$ can be dependent on $(p_f - p_i)/p_F$. This dependence comes from Eq. (2.12). As a result, we can obtain that at $T < T^*(B)$, the factor $d\rho/dT \propto A(B)T$ behaves as $A(B)T \propto T/(B - B_{c0})$, and at $T > T^*(B)$, it behaves as $A(B)T \propto 1/T$. These observations are in good agreement with the data obtained in measurements on $\text{YbRh}_2(\text{Si}_{0.95}\text{Ge}_{0.05})_2$ [102].

We note that, as in the case of the highly correlated liquid, the susceptibility $\chi(B, T)$ of the strongly correlated liquid is not a monotonic function of y and possesses a maximum as a function




FIG. 17: Magnetization $M(B)$ shown by filled squares as a function of magnetic field B [102]. The curve represents the field dependence of $M(B) = a_M \sqrt{B}$ given by Eq. (9.9) with a_M is a constnat.

of the temperature because the derivative $dM(B)/dB$ is the sum of two contributions. As it was shown in Section 5, the well-known empirical Kadowaki-Woods ratio [9], $K = A/\gamma_0^2 \simeq \text{const}$, is also obeyed in the case of the strongly correlated liquid. These results are in good agreement with facts [10, 102, 122].

As an application of the above consideration we study the $T - B$ phase diagram for the HF metal YbRh_2Si_2 [10] shown in Fig. 18. The LFL behavior is characterized by the effective mass $M^*(B)$ which diverges as $1/\sqrt{B - B_{c0}}$ [10]. We can conclude that Eq. (5.5) gives good description of this experimental fact, and $M^*(B)$ diverges at the point $B \rightarrow B_{c0}$ with $T_N(B = B_{c0}) = 0$. It is seen from Fig. 18, that the line separating the LFL state and NFL can be approximated by the function $c\sqrt{B - B_{c0}}$ with c being a parameter. Taking into account that the behavior of YbRh_2Si_2 strongly resembles the behavior of $\text{YbRh}_2(\text{Si}_{0.95}\text{Ge}_{0.05})_2$ [6, 102, 103, 122], we can conclude that in the NFL state the thermal expansion coefficient $\alpha(T)$ does not depend on T and the Grüneisen ratio as a function of temperature T diverges [6]. We are led to the conclusion that the entropy of the NFL state is given by Eq. (9.1). Taking into account that at relatively high temperatures the AF phase transition is of the second order [10], we predict that at lower temperatures this becomes the first order phase transition. Then, the described behavior of the Hall coefficient $R_H(B)$ is in good agreement with experimental facts [120].

Thus, we can conclude that the $T - B$ phase diagram of the heavy electron liquid with FC is in




FIG. 18: $T - B$ phase diagram for YbRh_2Si_2 [10, 102]. The T_N curve represents the field dependence of the Néel temperature. Line separating the antiferromagnetic (AF) and the non-Fermi liquid (NFL) state is a guide to the eye. The NFL state is characterized by the entropy S_{NFL} given by Eq. (9.1). Line separating the NFL state and LFL is $T^*(B - B_{c0}) = c\sqrt{B - B_{c0}}$, with c being an adjustable factor.

good agreement with the experimental $T - B$ phase diagram obtained in measurements on the HF metals YbRh_2Si_2 and $\text{YbRh}_2(\text{Si}_{0.95}\text{Ge}_{0.05})_2$.

9.2. Heavy fermion metals very near FCQPT on the ordered side

Let us consider the case when $\delta p_{FC} = (p_f - p_i)/p_F \ll 1$ and the electronic system of HF metal is very near FCQPT being on the ordered side. As we have seen in Section 5, Eq. (5.10), the application of magnetic field $(B - B_{c0})/B_{c0} \geq B_c$ removes the system from the ordered side placing it on the disordered one. As a result at $T \leq T_1(B)$, the effective mass $M^*(B)$ is given by Eqs. (8.5) and (8.11), while the corresponding resistivity is described in Section 8.2. In the absence of magnetic field or at $T_f \gg T > T_1(B)$, the system demonstrates the NFL behavior, the effective mass $M^*(T)$ is given by Eq. (2.12), and the entropy is given by Eq. (9.1). While the magnetic susceptibility $\chi(T) \propto M^*(T) \propto 1/T$, the thermal expansion coefficient $\alpha(T)$ is T -independent being determined by Eq. (9.3) and making the Grüneisen ratio divergent, as it follows from Eq. (9.5). Note, that the specific heat behaves as $C(T) \propto \sqrt{T}$ in both cases: when the electronic system is on the ordered side or on the disordered side of FCQPT, see Fig. 11 and Fig. 15. It follows from Eq. (2.12) that $\gamma(T) \propto T$. Therefore, the temperature dependent part $\Delta\rho(T)$ of the resistivity behaves

as $\Delta\rho(T) \propto \gamma(T) \propto T$. Thus, the system demonstrates the NFL regime, $\Delta\rho(T) \propto T$, when it is either in the highly correlated or in the strongly correlated regimes.

At some temperature T_c , the system can undergo the superconducting phase transition. In contrast to the conventional superconductors where the discontinuity $\delta C(T_c)$ in the specific heat at T_c is a linear function of T_c , $\delta C(T_c)$ is independent of the critical temperature T_c . As seen from Eqs. (3.23) and (3.24), both the discontinuity $\delta C(T_c)$ and the ratio $\delta C(T_c)/C_n(T_c)$ can be very large as compared to the conventional case [54, 123].

Recent experiments show that the electronic system of the HF metal CeCoIn₅ can be considered as the system located near FCQPT and containing FC. Indeed, under the application of magnetic field it behaves as the highly correlated liquid (see Section 8) with the effective mass $A(B) \propto (B - B_{c0})^{-4/3}$ [105], as it is seen from Eq. (8.16). While in the NFL regime, $\alpha(T) \propto \text{const}$ and the Grüneisen ratio diverges [124], see Eqs. (9.3) and (9.5), respectively. Estimations of δp_{FC} based on the evaluation of the magnetic susceptibility show that $\delta p_{FC} \simeq 0.044$ [123]. The obtained value of δp_{FC} allows to explain the relatively big value of the discontinuity $\delta C(T_c)$ [123] observed at $T_c = 2.3$ K in measurements on CeCoIn₅ [125]. Thus, we can conclude that $B_{cr} \simeq 0.003$, as it follows from Eq. (5.10), and the HF metal CeCoIn₅ can be considered as placed near FCQPT.

10. DISSYMMETRICAL TUNNELLING IN HF METALS AND HIGH- T_c SUPERCONDUCTORS

Experiments on the HF metals explore mainly their thermodynamic properties. It would be desirable to probe the other properties of the heavy electron liquid such as the probabilities of quasiparticle occupations, which are not directly linked to the density of states or to the behavior of the effective mass M^* [126]. Scanning tunnelling microscopy (STM) being sensitive to both the density of states and the probabilities of quasiparticle occupations is an ideal technique for studying such effects at quantum level.

The tunnelling current I through the point contact between two ordinary metals is proportional to the driving voltage V and to the squared modulus of the quantum mechanical transition amplitude t multiplied by the difference $N_1(0)N_2(0)(n_1(p, T) - n_2(p, T))$ [127]. Here $n(p, T)$ is the quasiparticle distribution function and $N(0)$ is the density of states of the corresponding metal. On the other hand, the wave function calculated in the WKB approximation and defining t is proportional to $(N_1(0)N_2(0))^{-1/2}$. As a result, the density of states is dropped out and the tunnelling current is

independent of $N_1(0)N_2(0)$. Upon taking into account that at $T \rightarrow 0$ the distribution $n(p, T \rightarrow 0) \rightarrow n_F(p)$, where $n_F(p)$ is the step function $\theta(p - p_F)$, one can check that within the LFL theory the differential tunnelling conductivity $\sigma_d(V) = dI/dV$ is a symmetric function of the voltage V .

In fact, the symmetry of $\sigma_d(V)$ holds provided that so called particle-hole symmetry is preserved as it is within the LFL theory. Therefore, the existence of the $\sigma_d(V)$ symmetry is quite obvious and common in the case of metal-to-metal contacts when these metals are ordinary metals and in their normal or superconducting states.

Now we turn to a consideration of the tunnelling current at low temperatures which in the case of ordinary metals is given by [127]

$$I(V) = 2|t|^2 \int [n_F(z - \mu) - n_F(z - \mu + V)] dz. \quad (10.1)$$

We use an atomic system of units: $e = m = \hbar = 1$, where e and m are electron charge and mass, respectively, and the energy z belongs to the interval E_0 given by Eq. (2.14)

$$\mu - 2T \leq z \leq \mu + 2T. \quad (10.2)$$

Since temperatures are low, we approximate the distribution function of ordinary metal by the step function n_F . It follows from Eq. (10.1) that quasiparticles with the energy z , $\mu - V \leq z \leq \mu$, contribute to the current, while $\sigma_d(V) \simeq 2|t|^2$ is a symmetrical function of V . In the case of the heavy electron liquid with FC, the tunnelling current is of the form [126]

$$I(V) = 2 \int [n_0(z - \mu) - n_F(z - \mu + V)] dz. \quad (10.3)$$

Here we have replaced the distribution function of ordinary metal by n_0 that is the solution of Eq. (2.8). We have also normalized the transition amplitude $|t|^2$ such that $|t|^2 = 1$. Assume that V satisfies the condition, $|V| \leq 2T$, while the current flows from the HF metal to the ordinary one. Quasiparticles of the energy z , $\mu - V \leq z$, contribute to $I(V)$, and the differential conductivity is given by the relation $\sigma_d(V) \simeq 2n_0(z \simeq \mu - V)$. If the sign of the voltage is changed, the direction of the current is also changed. In that case, quasiparticles of the energy z , $\mu + V \geq z$, contribute to $I(V)$, and the differential conductivity $\sigma_d(-V) \simeq 2(1 - n_0(z \simeq \mu + V))$. The dissymmetrical part $\Delta\sigma_d(V) = (\sigma_d(-V) - \sigma_d(V))$ of the differential conductivity is of the form

$$\Delta\sigma_d(V) \simeq 2[1 - (n_0(z - \mu \simeq V) - n_0(z - \mu \simeq -V))]. \quad (10.4)$$

It is worth noting that according to Eq. (10.4) we have $\Delta\sigma_d(V) = 0$ if the considered HF metal is replaced by an ordinary metal. Indeed, the effective mass is finite at $T \rightarrow 0$, then $n_0(T \rightarrow 0) \rightarrow n_F$

being given by Eq. (2.4), and $1 - n(z - \mu \simeq V) = n(z - \mu \simeq -V)$. One might say that the dissymmetrical part vanishes due to the particle-hole symmetry. On the other hand, there are no reasons to expect that $(1 - n_0(z - \mu \simeq V) - n_0(z - \mu \simeq -V)) = 0$. Thus, we conclude that the differential conductivity becomes a dissymmetrical function of the voltage.

To estimate $\Delta\sigma_d(V)$, we observe that it is zero when $V = 0$, because $n_0(p = p_F) = 1/2$ as it should be and it follows from Eq. (2.1) as well. It is seen from Eq. (10.4) that $\Delta\sigma_d(V)$ is an even function of both $(z - \mu)$ and V . Therefore we can assume that at low values of the voltage V the dissymmetrical part behaves as $\Delta\sigma_d(V) \propto V^2$. Then, the natural scale to measure the voltage is $2T$, as it is seen from Eq. (10.2). In fact, the dissymmetrical part is to be proportional to $(p_f - p_i)/p_F$. As a result, we obtain

$$\Delta\sigma_d(V) \simeq c \left(\frac{V}{2T} \right)^2 \frac{p_f - p_i}{p_F} \simeq c \left(\frac{V}{2T} \right)^2 \frac{S_0(r)}{x_{FC}}. \quad (10.5)$$

Here, $S_0(r)$ is the temperature independent part of the entropy (see Eq. (9.1)), c is a constant, which is expected to be of the order of a unit. This constant can be evaluated by using analytical solvable models. For example, calculations of c within a simple model, when the Landau functional $E[n(p)]$ is of the form [19]

$$E[n(p)] = \int \frac{p^2}{2M} \frac{d\mathbf{p}}{(2\pi)^3} + V_1 \int n(p)n(p) \frac{d\mathbf{p}}{(2\pi)^3}, \quad (10.6)$$

give $c \simeq 1/2$. It follows from Eq. (10.5), that when $V \simeq 2T$ and FC occupies a noticeable part of the Fermi volume, $(p_f - p_i)/p_F \simeq 1$, the dissymmetrical part becomes comparable with differential tunnelling conductivity, $\Delta\sigma_d(V) \sim V_d(V)$.

The dissymmetrical behavior of the tunnelling conductivity can be observed in measurements on both high- T_c metals in their normal state and the heavy fermion metals, for example, such as $\text{YbRh}_2(\text{Si}_{0.95}\text{Ge}_{0.05})_2$ or YbRh_2Si_2 which are expected to have undergone FCQPT. In the case of HF metals, upon the application of magnetic field B the effective mass is to diverge as given by Eq. (5.5). Here B_{c0} is the critical magnetic field which drives the HF metal to its magnetic field tuned quantum critical point. The value of the critical exponent $\alpha = -1/2$ is in good agreement with experimental observations collected on these metals [10, 102]. The measurements of $\Delta\sigma_d(V)$ have to be carried out applying magnetic field B_{c0} at temperatures $T^*(B) < T \leq T_f$. In the case of these metals, T_f is of the order of few Kelvin. We note that at sufficiently low temperatures, the application of magnetic field $B > B_{c0}$ leads to the restoration of the LFL behavior with $M^*(B)$ given by Eq. (5.5). As a result, the dissymmetrical behavior of the tunnelling conductivity vanishes [126].

The dissymmetrical differential conductivity $\Delta\sigma_d(V)$ can also be observed when both the high- T_c metal and the HF metal in question go from normal to superconducting. The reason is that $n_0(p)$

is again responsible for the dissymmetrical part of $\sigma_d(V)$. As we have seen in Section 3, this $n_0(p)$ is not appreciably disturbed by the pairing interaction which is relatively weak as compared to the Landau interaction forming the distribution function $n_0(p)$. In the case of superconductivity, we have to take into account that the ratio,

$$\frac{N_s(E)}{N(0)} = \frac{|E|}{\sqrt{E^2 - \Delta^2}}, \quad (10.7)$$

comes into the play because the density of states $N_s(E)$ of the superconducting metal is zero in the gap, that is when $|E| \leq |\Delta|$. Here E is the quasiparticle energy, while the normal state quasiparticle energy is $\varepsilon - \mu = \sqrt{E^2 - \Delta^2}$. Now we can adjust Eq. (10.4) for the case of superconducting HF metal, multiplying the right hand side of Eq. (10.4) by $N_s/N(0)$ and replacing the quasiparticle energy $z - \mu$ by $\sqrt{E^2 - \Delta^2}$ with E being represented by the voltage V . As a result, Eq. (10.5) can be presented in the following form

$$\Delta\sigma_d(V) \simeq \left| \frac{V}{\Delta} \right| \frac{(\sqrt{V^2 - \Delta^2})^2}{|\Delta|\sqrt{V^2 - \Delta^2}} \frac{p_f - p_i}{p_F} \simeq \sqrt{1 - \left[\frac{\Delta}{V} \right]^2} \left[\frac{V}{\Delta} \right]^2 \frac{S_0(r)}{x_{FC}}. \quad (10.8)$$

It is seen from Eqs. (10.5) and (10.8) that, as in the case described by Eq. (9.7) when the abrupt change in the Hall coefficient is defined by $S_0(r)$, the dissymmetrical part of the differential tunnelling conductivity is also proportional to the $S_0(r)$ vanishing at $r \rightarrow 0$. Note that the scale $2T$ entering Eq. (10.5) is replaced by the scale Δ in Eq. (10.8). In the same way, as Eq. (10.5) is valid up to $V \sim 2T$, Eq. (10.8) is valid up to $V \sim 2|\Delta|$. It is seen from Eq. (10.8) that the dissymmetrical part of the differential tunnelling conductivity becomes as large as the differential tunnelling conductivity at $V \sim 2|\Delta|$ provided that FC occupies a large part of the Fermi volume, $(p_f - p_i)/p_F \simeq 1$. In the case of a d -wave gap, the right hand side of Eq. (10.8) has to be integrated over the gap distribution. As a result, $\Delta\sigma_d(V)$ is expected to be finite even at $V = \Delta_1$, where Δ_1 is the maximum value of the d -wave gap [126].

The presence of an electronic inhomogeneity in $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_{8+x}$ was recently discovered in observations using scanning tunnelling microscopy and spectroscopy [128]. This inhomogeneity is manifested as spatial variations in the local density of states (LDOS) spectrum, in the low-energy spectral weight, and in the magnitude of the superconducting energy gap. The inhomogeneity observed in the integrated LDOS is not induced by impurities, but rather is intrinsic in nature. The observations allowed to relate the magnitude of the integrated LDOS to the local oxygen doping concentration [128]. Spatial variation of the tunnelling differential conductance spectrum are shown in Fig. 19. The curves 1 and 2 can be considered as corresponding to the normal state of high- T_c superconductor.




FIG. 19: Spatial variation of the tunnelling differential conductance spectra measured on $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_{8+x}$. Curves 1 and 2 are taken at positions where the integrated LDOS is very small. The low differential conductance and the absence of a superconducting gap are indicative of insulating behavior. Curve 3 is for a large gap of 65 meV, with low coherence peaks. The integrated value of the LDOS at the position for curve 3 is small but larger than those in curves 1 and 2. Curve 4 is for a gap of 40 meV, which is close to the mean value of the gap distribution. Curve 5, taken at the position with the highest integrated LDOS, is for the smallest gap of 25 meV with two very sharp coherence peaks [128].

The other curves can be viewed as corresponding to high- T_c superconductors with different oxygen doping concentrations. It is seen from Fig. 19 that the tunnelling differential conductivity is strongly dissymmetrical in both the normal state and the superconducting state of the $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_{8+x}$ compound. Therefore, we can conclude that the dissymmetrical tunnelling described by Eqs. (10.5) and (10.8) is in good qualitative agreement with the experimental facts displayed in Fig. 19.

The dissymmetrical tunnelling conductivity given by Eqs. (10.5) and (10.8) can be observed in measurements on the heavy fermion metal CeCoIn_5 in its superconducting state and its NFL normal state. As it was discussed in Subsection 9.2, this metal is expected to have undergone FCQPT.

11. SUMMARY AND CONCLUSION

Through out this paper we have discussed the manifestations of the fermion condensation, which can be compared to the Bose-Einstein condensation. A number of experimental evidences have been

presented that are supportive to the idea of the existence of FC in different natural and artificial substances. We have demonstrated also that numerous experimental facts collected in a whole variety of materials, belonging to the high- T_c superconductors, heavy fermion metals and strongly correlated 2D structures, can be explained within the framework of the theory based on FCQPT.

We have shown that FCQPT separates the regions of LFL and strongly correlated liquids. Beyond the FCQPT point the quasiparticle system is divided into two subsystems, one containing normal quasiparticles, while the other being occupied by FC localized at the Fermi level. In the superconducting state the quasiparticle dispersion $\varepsilon(\mathbf{p})$ in systems with FC can be approximated by two straight lines, characterized by effective masses M_{FC}^* and M_L^* , and intersecting near the binding energy E_0 which is of the order of the superconducting gap. The same quasiparticle picture and the energy scale E_0 persist in the normal state.

We have demonstrated that fermion systems with FC have features of "quantum protectorate" and shown that the theory of high- T_c superconductivity, based on FCQPT and on the conventional theory of superconductivity, permits the description of high values of T_c and of the maximum value of the gap Δ_1 , which may be as big as $\Delta_1 \sim 0.1\varepsilon_F$ or even larger. We have also traced the transition from conventional superconductors to high- T_c ones. We have shown by a simple, although self-consistent analysis that both the pseudogap state and the general features of the shape of the critical temperature $T_c(x)$ as a function of the number density x of the mobile carriers in the high- T_c compounds can be understood within the framework of the theory.

We have also shown that striking experimental results on the magnetic-field induced LFL in high- T_c metals, which unveil the nature of the high- T_c superconductivity, suggest that FCQPT and the emergence of the novel quasiparticles with effective mass strongly depending on the magnetic field and temperature and resembling the Landau quasiparticles are qualities intrinsic to the electronic system of the high- T_c superconductors.

We have provided explanations of the experimental data on the divergence of the effective mass in 2D electron liquid and in 2D ^3He , as well as shown that above the critical point of FCQPT the system exhibits the LFL behavior. The behavior of the heavy electron liquid approaching FCQPT from the disordered phase can be viewed as the highly correlated one because the effective mass is very large and strongly depends on the density, temperature and magnetic fields.

At different temperatures, the behavior in magnetic fields of the highly correlated electron liquid approaching FCQPT from the disordered phase has been considered. We have shown that at sufficiently high temperatures $T_1(B) < T$, the effective mass starts to depend on T , $M^* \propto T^{-1/2}$.

This $T^{-1/2}$ dependence of the effective mass at elevated temperatures leads to the non-Fermi liquid behavior of the resistivity, $\rho(T) \propto T$. The application of magnetic field B restores the common T^2 behavior of the resistivity. If the magnetic field $(B - B_{c0})$ decreases to zero, the effective mass M^* diverges as $M^* \propto (B - B_{c0})^{-2/3}$. At finite magnetic fields, the regime NFL is restored at some temperature $T_1(B) \propto (B - B_{c0})$ with $M^*(T) \propto T^{-2/3}$. We have demonstrated that this $B - T$ phase diagram has a strong impact on MR of the highly correlated electron liquid. At fixed B , MR as a function of the temperature exhibits a transition from the negative values of MR at $T \rightarrow 0$ to the positive values at $T \propto (B - B_{c0})$. While at low temperatures and elevated magnetic fields, MR goes from positive to negative. This behavior was observed in the heavy fermion metals.

We have demonstrated that the strongly correlated electron liquid with FC, which exhibits strong deviations from the LFL behavior down to lowest temperatures, can be driven into LFL by applying a magnetic field B . If the magnetic field $(B - B_{c0})$ decreases to zero, the effective mass M^* diverges as $M^* \propto 1/\sqrt{B - B_{c0}}$ and the Néel temperature T_N of the AF phase transition tends to zero, $T_N(B \rightarrow B_{c0}) \rightarrow 0$. The NFL regime is restored at some temperature $T^*(B) \propto \sqrt{B - B_{c0}}$. In that case and at $T \rightarrow 0$ and $B = B_{c0}$, the Grüneisen ratio as a function of temperature T diverges. While the entropy $S(T)$ possesses the specific low temperature behavior, $S(T) \propto S_0 + a\sqrt{T} + bT$ with S_0 , a and b are temperature independent constants. We have shown that the obtained $T - B$ phase diagram is in good agreement with the experimental $T - B$ phase diagram obtained in measurements on the HF metals YbRh_2Si_2 and $\text{YbRh}_2(\text{Si}_{0.95}\text{Ge}_{0.05})_2$. We have also demonstrated that the abrupt jump in the Hall coefficient $R_H(B \rightarrow B_{c0}, T \rightarrow 0)$ is determined by the presence of FC. We have observed that the second order AF phase transition changes to the first order one below $T_{crit} < T_N(B = 0)$ making the corresponding quantum and thermal critical fluctuations vanish at the jump. Therefore, the abrupt jump and the divergence of the effective mass taking place at $T_N \rightarrow 0$ are defined by the behavior of quasiparticles rather than by the corresponding thermal and quantum critical fluctuations.

We have predicted that the differential tunnelling conductivity between a metallic point and an ordinary metal, which is commonly symmetric as a function of the voltage, becomes noticeably dissymmetrical when the ordinary metal is replaced by a HF metal, the electronic system of which has undergone FCQPT. This dissymmetry can be observed when the HF metal is both normal and superconducting. We have also discussed possible experiments to study the dissymmetry in measurements on the HF metals. In the case of the high- T_c superconductors, our consideration is in good agreement with available data.

In conclusion, we have shown that in contrast to conventional quantum phase transitions, whose

physics is dominated by thermal and quantum fluctuations and characterized by the absence of quasiparticles, the physics of Fermi systems and the heavy electron liquid near FCQPT or undergone FCQPT is determined by quasiparticles resembling the Landau quasiparticles. We have shown that the Landau paradigm based on the notion of quasiparticles and order parameters is still applicable when considering the low temperature properties of the heavy electron liquid, whose understanding has been problematic largely because of the absence of theoretical guidance. In contrast with the conventional Landau quasiparticles, the effective mass of the considered quasiparticles strongly depends on the temperature, applied magnetic fields, the number density x , pressure, etc. These quasiparticles and the order parameter are well defined and capable of describing both the LFL and the NFL behaviors of both the high- T_c superconductors and the HF metals and their universal thermodynamic properties down to the lowest temperatures. This system of quasiparticles determines the recovery of the LFL behavior under applied magnetic fields and preserves the Kadowaki-Woods ratio. Thus, we obtain a unique possibility to control the essence of HF systems and accordingly of the HF metals by magnetic fields in a wide range of temperatures.

Finally, our general consideration suggests that FCQPT and the emergence of novel quasiparticles at QCP and behind QCP and resembling the Landau quasiparticles are qualities intrinsic to strongly correlated substances, while FCQPT can be viewed as the universal cause of the non-Fermi liquid behavior observed in different metals and liquids.

12. ACKNOWLEDGMENTS

We are grateful to P. Coleman, V.A. Khodel and M. Norman for valuable discussions.

This work was supported in part by the Russian Foundation for Basic Research, project No. 01-02-17189 and in part by INTAS, project No. 03-51-6170. The visit of VRS to Clark Atlanta University has been supported by NSF through a grant to CTSPS. VRS is grateful to the Racah Institute of Physics for the hospitality during his stay at the Hebrew University, Jerusalem. MYaA is grateful to the Binational Science Foundation, grant 2002064 and Israeli Science Foundation, grant 174/03. AZM is supported by US DOE, Division of Chemical Sciences, Office of Basic Energy Sciences, Office of Energy Research. PKG is supported by the grants of Russian Academy of Sciences.

[1] G.R. Stewart, Rev. Mod. Phys. **73**, 797 (2001).

- [2] C.M. Varma, Z. Nussionov, and W. van Saarloos, Phys. Rep. **361**, 267 (2002).
- [3] S. Sachdev, *Quantum Phase transitions* (Cambridge, Cambridge University Press, 1999).
- [4] M. Vojta, Rep. Prog. Phys. **66**, 2069 (2003).
- [5] D.V. Shopova and D.I. Uzunov, Phys. Rep. **379**, 1 (2003).
- [6] R. K  chler, N. Oeschler, P. Gegenwart, T. Cichorek, K. Neumaier, O. Tegus, C. Geibel, J. A. Mydosh, F. Steglich, L. Zhu, and Q. Si, Phys. Rev. Lett. **91**, 066405 (2003).
- [7] M.Ya. Amusia, A.Z. Msezane, and V.R. Shaginyan, Phys. Lett. A **320**, 459 (2004).
- [8] V.R. Shaginyan, JETP Lett. **79**, 286 (2004).
- [9] K. Kadowaki and S.B. Woods, Solid State Commun. **58**, 507 (1986).
- [10] P. Gegenwart, J. Custers, C. Geibel, K. Neumaier, T. Tayama, K. Tenya, O. Trovarelli, and F. Steglich, Phys. Rev. Lett. **89**, 056402 (2002).
- [11] C. Proust, E. Boaknin, R. W. Hill, L. Taillefer, and A. P. Mackenzie, Phys. Rev. Lett. **89**, 147003 (2002).
- [12] A.J. Millis, A.J. Schofield, G.G. Lonzarich, and S.A. Grigera, Phys. Rev. Lett. **88**, 217204 (2002).
- [13] A. Bianchi, R. Movshovich, I. Vekhter, P. G. Pagliuso, and J. L. Sarrao, Phys. Rev. Lett. **91**, 257001 (2003); F. Ronning, C. Capan, A. Bianchi, R. Movshovich, A. Lacerda, M. F. Hundley, J. D. Thompson, P. G. Pagliuso, and J. L. Sarrao, Phys. Rev. B **71**, 104528 (2005).
- [14] J. Paglione, M.A. Tanatar, D.G. Hawthorn, E. Boaknin, F. Ronning, R.W. Hill, M. Sutherland, L. Taillefer, C. Petrovic, and P.C. Canfield, cond-mat/0405157.
- [15] S.T. Belyaev, Sov. Phys. JETP, **7**, 289 (1958).
- [16] S.T. Belyaev, Sov. Phys. JETP, **7**, 299 (1958).
- [17] V.A. Khodel and V.R. Shaginyan, JETP Lett. **51**, 553 (1990).;
- [18] V.A. Khodel and V.R. Shaginyan, Nucl. Phys. A **555**, 33 (1993).
- [19] V.A. Khodel, V.R. Shaginyan, and V.V. Khodel, Phys. Rep. **249**, 1 (1994).
- [20] J. Dukelsky, V.A. Khodel, P. Schuck, and V.R. Shaginyan, Z. Phys. **102**, 245 (1997); V.A. Khodel and V.R. Shaginyan, Condensed Matter Theories, **12**, 222 (1997).
- [21] V.R. Shaginyan, Phys. Lett. A **249**, 237 (1998).
- [22] G. E. Volovik, JETP Lett. **53**, 222 (1991).
- [23] I.E. Dzyaloshinskii, J. Phys. I (France) **6**, 119 (1996).
- [24] D. Lidsky, J. Shiraishi, Y. Hatsugai, and M. Kohmoto, Phys. Rev. B **57**, 1340 (1998).
- [25] V.Yu. Irkhin, A.A. Katanin, and M.I. Katsnelson, Phys. Rev. Lett. **89**, 076401 (2002).

- [26] V.R. Shaginyan, JETP Lett. **77**, 178 (2003).
- [27] E.M. Lifshitz and L.P. Pitaevskii, *Statistical Physics* (Part 2, Butterworth-Heinemann, Oxford, 1999).
- [28] L.Ja. Pomeranchuk, JETP **35**, 524 (1958).
- [29] M.Ya. Amusia and V.R. Shaginyan, JETP Lett. **73**, 232 (2001);
- [30] M.Ya. Amusia and V.R. Shaginyan, Phys. Rev. B **63**, 224507 (2001); V.R. Shaginyan, Physica B **312-313C**, 413 (2002).
- [31] V.R. Shaginyan, JETP Lett. **77**, 99 (2003).
- [32] V.M.Yakovenko and V.A. Khodel, JETP Lett. **78**, 398 (2003); cond-mat/0308380.
- [33] A. Casey, H. Patel, J. Nye'ki, B. P. Cowan, and J. Saunders, Phys. Rev. Lett. **90**, 115301 (2003).
- [34] A.A. Shashkin, S.V. Kravchenko, V.T. Dolgoplov, and T.M. Klapwijk, Phys. Rev. B **66**, 073303 (2002); A.A. Shashkin, M. Rahimi, S. Anissimova, S.V. Kravchenko V.T. Dolgoplov, and T. M. Klapwijk, Phys. Rev. Lett. **91**, 046403 (2003).
- [35] J. Boronat, J. Casulleras, V. Grau, E. Krotscheck, and J. Springer, Phys. Rev. Lett. **91**, 085302 (2003).
- [36] Y. Zhang, V. M. Yakovenko, and S. Das Sarma, Phys. Rev. B **71**, 115105 (2005).
- [37] Y. Zhang and S. Das Sarma, Phys. Rev. B **70**, 035104 (2004).
- [38] V.A. Khodel, V.R. Shaginyan, and M.V. Zverev, JETP Lett. **65**, 253 (1997).
- [39] V.R. Shaginyan, J.G. Han, and J. Lee, Phys. Lett. A **329**, 108 (2004).
- [40] R.B. Laughlin and D. Pines, Proc. Natl. Acad. Sci. USA **97**, 28 (2000).
- [41] P.W. Anderson, cond-mat/0007185; cond-mat/0007287.
- [42] V.A. Khodel, J.W. Clark, and V.R. Shaginyan, Solid Stat. Comm. **96**, 353 (1995).
- [43] S.A. Artamonov and V.R. Shaginyan, JETP **92**, 287 (2001).
- [44] P. V. Bogdanov, A. Lanzara, S. A. Kellar, X. J. Zhou, E. D. Lu, W. J. Zheng, G. Gu, J.-I. Shimoyama, K. Kishio, H. Ikeda, R. Yoshizaki, Z. Hussain, and Z. X. Shen, Phys. Rev. Lett. **85**, 2581 (2000).
- [45] A. Kaminski, M. Randeria, J. C. Campuzano, M. R. Norman, H. Fretwell, J. Mesot, T. Sato, T. Takahashi, and K. Kadowaki, Phys. Rev. Lett. **86**, 1070 (2001).
- [46] T. Valla, A. V. Fedorov, P. D. Johnson, B. O. Wells, S. L. Hulbert, Q. Li, G. D. Gu, and N. Koshizuka, Science **285**, 2110 (1999);
T. Valla, A. V. Fedorov, P. D. Johnson, Q. Li, G. D. Gu, and N. Koshizuka, Phys. Rev. Lett. **85**, 828 (2000).
- [47] G. Grüner, *Density Waves in Solids* (Addison-Wesley, Reading, MA, 1994).
- [48] L. Tassini, F. Venturini, Q.-M. Zhang, R. Hackl, N. Kikugawa, and T. Fujita, Phys. Rev. Lett. **95**,

- 117002 (2005).
- [49] J. Bardeen, L.N. Cooper, and J.R. Schrieffer, Phys. Rev. **108**, 1175 (1957).
 - [50] D.R. Tilley and J. Tilley, *Superfluidity and Superconductivity*, (Bristol, Hilger, 1985).
 - [51] N.N. Bogoliubov, Nuovo Cimento **7**, 794 (1958).
 - [52] M.Ya. Amusia and V.R. Shaginyan, JETP Lett. **77**, 671 (2003).
 - [53] H. Matsui, T. Sato, T. Takahashi, S.-C. Wang, H.-B. Yang, H. Ding, T. Fujii, T. Watanabe, and A. Matsuda, Phys. Rev. Lett. **90**, 217002 (2003).
 - [54] M.Ya. Amusia, S.A. Artamonov, and V.R. Shaginyan, JETP Lett. **74**, 435 (2001).
 - [55] A. Paramekanti, M. Randeria, and N. Trivedi, Phys. Rev. Lett. **87**, 217002 (2001); A. Paramekanti, M. Randeria, and N. Trivedi, cond-mat/0305611.
 - [56] P. W. Anderson, P. A. Lee, M. Randeria, T. M. Rice, N. Trivedi, and F. C. Zhang, J Phys. Condens. Matter **16**, R755 (2004).
 - [57] V.R. Shaginyan, JETP Lett. **68**, 527 (1998).
 - [58] M.Ya. Amusia and V.R. Shaginyan, Phys. Lett. A **298**, 193 (2002).
 - [59] M. Kugler, M. Fischer, Ch. Renner, S. Ono, and Y. Ando, Phys. Rev. Lett. **86**, 4911 (2001).
 - [60] A.A. Abrikosov, Phys. Rev. B **52**, R15738 (1995); A.A. Abrikosov, cond-mat/9912394.
 - [61] N.-C. Yeh, C.-T. Chen, G. Hammer, J. Mannhart, A. Schmeh, C. W. Schneider, R. R. Schulz, S. Tajima, K. Yoshida, D. Garrigus, and M. Strasik, Phys. Rev. Lett. **87**, 087003 (2001).
 - [62] A. Biswas, P. Fournier, M. M. Qazilbash, V. N. Smolyaninova, H. Balci, and R. L. Greene, Phys. Rev. Lett. **88**, 207004 (2002).
 - [63] J.A. Skinta, M.-S. Kim, T.R. Lemberger, T. Greibe, and M. Naito, Phys. Rev. Lett. **88**, 207005 (2002).
 - [64] J.A. Skinta, T.R. Lemberger, T. Greibe, and M. Naito, Phys. Rev. Lett. **88**, 207003 (2002).
 - [65] C.-T. Chen, P. Seneor, N.-C. Yeh, R.P. Vasquez, L.D. Bell, C.U. Jung, J.Y. Kim, M.-S. Park, H.-J. Kim, and S.-I. Lee, Phys. Rev. Lett. **88**, 227002 (2002).
 - [66] N. Metoki, Y. Haga, Y. Koike, and Y. Onuki, Phys. Rev. Lett. **80**, 5417 (1998).
 - [67] T. Honma, Y. Haga, and E. Yamamoto, J. Phys. Soc. Jpn. **68**, 338 (1999).
 - [68] M.Ya. Amusia and V.R. Shaginyan, JETP Lett. **76**, 651 (2002).
 - [69] N. Miyakawa, J. F. Zasadzinski, L. Ozyuzer, P. Guptasarma, D. G. Hinks, C. Kendziora, and K. E. Gray, Phys. Rev. Lett. **83**, 1018 (1999).
 - [70] C.M. Varma, P.B. Littlewood, S. Schmitt-Rink, E. Abrahams, and A.E. Ruckenstein, Phys. Rev. Lett. **63**, 1996 (1989).

- [71] C.M. Varma, P.B. Littlewood, S. Schmitt-Rink, E. Abrahams, and A.E. Ruckenstein, Phys. Rev. Lett. **64**, 497 (1990).
- [72] A. Ino, C. Kim, M. Nakamura, T. Yoshida, T. Mizokawa, A. Fujimori, Z.-X. Shen, T. Kakeshita, H. Eisaki, and S. Uchida, Phys. Rev. B **65**, 094504 (2002).
- [73] X.J. Zhou, T. Yoshida, A. Lanzara, P.V. Bogdanov, S.A. Kellar, K.M. Shen, W. L. Yang, F. Ronning, T. Sasagawa, T. Kakeshita, T. Noda, H. Eisaki, S. Uchida, C. T. Lin, F. Zhou, J.W. Xiong, W.X. Ti, Z.X. Zhao, A. Fujimori, Z. Hussain, and Z.-X. Shen, Nature **423**, 398 (2003).
- [74] W. J. Padilla, Y. S. Lee, M. Dumm, G. Blumberg, S. Ono, K. Segawa, S. Komiya, Y. Ando, D. N. Basov, Phys. Rev. B **72**, 060511 (2005).
- [75] T. Valla, A.V. Fedorov, P.D. Johnson, and S.L. Hulbert, Phys. Rev. Lett. **83**, 2085 (1999).
- [76] D.L. Feng, A. Damascelli, K.M. Shen, N. Motoyama, D.H. Lu, H. Eisaki, K. Shimizu, J.-I. Shimoyama, K. Kishio, N. Kaneko, M. Greven, G. D. Gu, X. J. Zhou, C. Kim, F. Ronning, N.P. Armitage, and Z.-X. Shen, Phys. Rev. Lett. **88**, 107001 (2002).
- [77] D.S. Dessau, B O. Wells, Z.X. Shen, W. E. Spicer, A.J. Arko, R.S. List, D. B. Mitzi, and A. Kapitulnik, Phys. Rev. Lett. **66**, 2160 (1991).
- [78] A.B. Migdal, *Theory of Finite Fermi Systems and Applications to Atomic Nuclei* (Benjamin, Reading, MA, 1977).
- [79] Z.M. Yusof, B.O. Wells, T. Valla, A.V. Fedorov, P.D. Johnson, Q. Li, C. Kendziora, S. Jian, and D. G. Hinks, Phys. Rev. Lett. **88**, 167006 (2002).
- [80] S. Nakamae, K. Behnia, N. Mangkorntong, M. Nohara, H. Takagi, S. Yates, and N.E. Hussey, Phys. Rev. B **68**, 100502 (2003).
- [81] N.E. Hussey, M. Abdel-Jawad, A. Carrington, A.P. Mackenzie, and L. Balicas, Nature **425**, 814 (2003).
- [82] S. A. Artamonov, V.R. Shaginyan, and Yu.G. Pogorelov, JETP Lett. **68**, 942 (1998).
- [83] Yu.G. Pogorelov and V.R. Shaginyan, JETP Lett. **76**, 532 (2002).
- [84] M. de Llano and J. P. Vary, Phys. Rev. C **19**, 1083 (1979); M. de Llano, A. Plastino, and J.G. Zabolitsky, Phys. Rev. C **20**, 2418 (1979).
- [85] M.V. Zverev and M. Baldo, J. Phys. Condens. Matter **11**, 2059 (1999).
- [86] V.R. Shaginyan, M.Z. Msezane, and M.Ya. Amusia, Phys. Lett. A **338**, 393 (2005).
- [87] D. Takahashi, S. Abe, H. Mizuno, D. A. Tayurskii, K. Matsumoto, H. Suzuki, and Y. Onuki, Phys. Rev. B **67**, 180407 (2003).
- [88] V.A. Khodel and P. Schuck, Z. Phys. B **104**, 505 (1997).

- [89] N. Tsujii, H. Kontani, and K. Yoshimura, Phys. Rev. Lett. **94**, 057201 (2005).
- [90] R. Bel, K. Behnia, C. Proust, P. van der Linden, D.K. Maude, and S.I. Vedenev, Phys. Rev. Lett. **92**, 17703 (2004).
- [91] J. Korringa, Physica (Utrecht) **16**, 601 (1950).
- [92] G.-q. Zheng, T. Sato, Y. Kitaoka, M. Fujita, and K. Yamada, Phys. Rev. Lett. **90**, 197005 (2003).
- [93] C.L. Kane and M.P.A. Fisher, Phys. Rev. Lett. **76**, 3192 (1996).
- [94] T. Senthil and M.P.A. Fisher, Phys. Rev. B **62**, 7850 (2000).
- [95] A. Houghton, S. Lee, and J.P. Marston, Phys. Rev. B **65**, 22503 (2002).
- [96] A.P. Mackenzie, S.R. Julian, D.C. Sinclair, and C. T. Lin, Phys. Rev. B **53**, 5848 (1996).
- [97] M.Ya. Amusia and V.R. Shaginyan, Phys. Lett. A **315**, 288 (2003).
- [98] K.D. Morhard, C. Bäuerle, J. Bossy, Yu. Bunkov, S.N. Fisher, and H. Godfrin, Phys. Rev. B **53**, 2658 (1996).
- [99] A. Casey, H. Patel, J. Nyéki, B. P. Cowan, and J. Saunders, Phys. Rev. Lett. **90**, 115301 (2003).
- [100] M. Pfizner and P. Wölfe, Phys. Rev. B **33**, 2003 (1986).
- [101] S.V. Kravchenko and M.P. Sarachik, Rep. Prog. Phys. **67**, 1 (2004).
- [102] J. Custers, P. Gegenwart, H. Wilhelm, K. Neumaier, Y. Tokiwa, O. Trovarelli, C. Geibel, F. Steglich, C. Pépin, and P. Coleman, Nature **424**, 524 (2003).
- [103] C. Pépin, Phys. Rev. Lett. **94**, 066402 (2005).
- [104] S.L. Bud'ko, E. Morosan, and P.C. Canfield, Phys. Rev. **69**, 014415 (2004).
- [105] J. Paglione, M.A. Tanatar, D.G. Hawthorn, E. Boaknin, R.W. Hill, F. Ronning, M. Sutherland, and L. Taillefer, Phys. Rev. Lett. **91**, 246405 (2003).
- [106] T. Senthil, M. Vojta, and S. Sachdev, Phys. Rev. B **69**, 035111 (2004); T. Senthil, S. Sachdev, and M. Vojta, Physica B **359-361**, 9 (2005).
- [107] T. Senthil, A. Vishwanath, L. Balents, S. Sachdev, and M.P.A. Fisher, Science **303**, 1490 (2004).
- [108] R. Küchler, P. Gegenwart, K. Heuser, E.-W. Scheidt, G. R. Stewart, and F. Steglich, Phys. Rev. Lett. **93**, 096402 (2004).
- [109] A. Bianchi, R. Movshovich, I. Vekhter, P.G. Pagliuso, and J.L. Sarrao, Phys. Rev. Lett. **91**, 257001 (2003).
- [110] L. Zhu, M. Garst, A. Rosch, and Q. Si, Phys. Rev. Lett. **91**, 066404 (2003).
- [111] K. Ishida, K. Okamoto, Y. Kawasaki, Y. Kitaoka, O. Trovarelli, C. Geibel, and F. Steglich, Phys. Rev. Lett. **89**, 107202 (2002).

- [112] P. Coleman, *Lectures on the Physics of Highly Correlated Electron Systems VI* (Editor F. Mancini, American Institute of Physics, New York, 2002, p. 79).
- [113] P. Coleman and A.J. Schofield, *Nature* **433**, 226 (2005).
- [114] J.W. Clark, V.A. Khodel, and M.V. Zverev, *Phys. Rev B* **71**, 012401 (2005).
- [115] E.M. Lifshitz and L.P. Pitaevskii, *Statistical Physics* (Part 1, Butterworth-Heinemann, 2000, p. 168).
- [116] V.R. Shaginyan, *JETP Lett.* **80**, 263 (2004).
- [117] J.W. Clark, V.A. Khodel, M.V. Zverev, and V.M. Yakovenko, *Phys. Rep.* **391**, 123 (2004).
- [118] M.V. Zverev, V.A. Khodel, and V.R. Shaginyan, *JETP Lett.* **65**, 863 (1997).
- [119] P. Nozières, *J. Phys. I (France)* **2**, 443 (1992).
- [120] S. Paschen, T. Lühmann, S. Wirth, P. Gegenwart, O. Trovarelli, C. Geibel, F. Steglich, P. Coleman and Q. Si, *Nature* **432**, 881 (2004).
- [121] V.R. Shaginyan, K.G. Popov, and S.A. Artamonov, *JETP Lett.* **82**, 234 (2005).
- [122] P. Gegenwart, J. Custers, Y. Tokiwa, C. Geibel, and F. Steglich, *Phys. Rev. Lett.* **94**, 076402 (2005).
- [123] V.A. Khodel, M.V. Zverev, and V.M. Yakovenko, cond-mat/0508275.
- [124] N. Oeschler, P. Gegenwart, M. Lang, R. Movshovich, J.L. Sarrao, J.D. Thompson, and F. Steglich, *Phys. Rev. Lett.* **91**, 076402 (2003).
- [125] E. D. Bauer, J.D. Thompson, J.L. Sarrao, L.A. Morales, F. Wastin, J. Rebizant, J.C. Griveau, P. Javorsky, P. Boulet, E. Colineau, G. H. Lander, and G. R. Stewart, *Phys. Rev. Lett.* **93**, 147005 (2004).
- [126] V.R. Shaginyan, *JETP Lett.* **81**, 222 (2005).
- [127] A.M. Zagoskin, *Quantum Theory of Many-Body Systems* (Springer-Verlag New York, Inc., 1998).
- [128] S.H. Pan, J.P. O'Neal, R.L. Badzey, C. Chamon, H. Ding, J.R. Engelbrecht, Z. Wang, H. Eisaki, S. Uchida, A.K. Gupta, K.-W. Ng, E.W. Hudson, K.M. Lang, and J. C. Davis, *Nature* **413**, 282 (2001).